

15th International Conference on X-Ray Lasers 2016

Conference Program and Book of Abstracts

May 22nd - 27th, 2016

Nara Kasugano International Forum, Iraka

*The 15th International Conference on X-ray Lasers
May 22-27, 2016, Nara Kasugano International Forum*

Conference co-chairs:

T. Kawachi, S. V. Bulanov, H. Daido, and Y. Kato

<http://www.kansai.qst.go.jp/icxrl2016/>

Organized by:

KPSI/QST

Kansai Photon Science Institute,
National Institutes for Quantum and
Radiological Science and Technology
Japan Atomic Energy Agency
GPI

The Graduate School for the
Creation of New Photonics Industries

Topics:

Laser pumped and discharge pumped X-ray Lasers
Injection/Seeding of X-ray Amplifiers
Higher order harmonics generation
X-ray free electron lasers
Novel schemes for coherent XUV, X-ray and γ -ray generation
X-rays and γ -rays for fundamental science
Integration of X-ray lasers, XFELs and super-intense lasers
Applications and industrial uses
X-ray imaging and optics



15th International Conference on X-Ray Lasers 2016

Conference Co-chairs

Tetsuya Kawachi (National Institute for Quantum and Radiological Science and Technology)
Sergei Bulanov (National Institute for Quantum and Radiological Science and Technology)
Hiroyuki Daido (Japan Atomic Energy Agency)
Yoshiaki Kato (The Graduate School for the Creation of New Photonics Industries)

International Advisory Board

L. M. Chen (IOP, Beijing, China)
P. Corkum (NRC/Univ. of Ottawa, Canada)
J. Feldhauss (Desy, Germany)
K. A. Janulewicz (WAT, Poland)
A. Klisnick (ISMO, France)
O. Kocharovskaya (Texas A&M Univ., USA)
M. Kozlova (IOP, Czech Republic)
C. L. S. Lewis (Queen's Univ. Belfast, UK)
C. H. Nam (IBS-CoReLS, Korea)
P. V. Nickles (MBI, Germany)
J. Nilsen (LLNL, USA)
J. J. Rocca (Colorado State Univ., USA)
S. Sebban (LOA, France/ELI, Czech Republic)
S. Suckewer (Princeton Univ., USA)
G. J. Tallents (Univ. of York, UK)
A. Vinogradov (LPI, Russia)

Local Organizing Committee

Masataka Kado (QST) (Chair)
Shinsuke Fujioka (ILE, Osaka Univ.)
Takeshi Higashiguchi (Utsunomiya Univ.)
Ryosuke Kodama (IAI, Osaka Univ.)
Shoichi Kubodera (Soka Univ.)
Katsumi Midorikawa (RIKEN)
Shuji Miyamoto (Univ. Hyogo)
Hakaru Mizoguchi (Gigaphoton)
Atsushi Momose (Tohoku Univ.)
Shinichi Namba (Hiroshima Univ.)
Makina Yabashi (RIKEN)
Hitoki Yoneda (Univ. Electro-Commun.)
Timur Zh. Esirkepov (QST)
Masaki Kando (QST)
Hiromitsu Kiriya (QST)
Kiminori Kondo (QST)
Akira Sasaki (QST)

Supporting Institutions

Kansai Photon Science Institute (KPSI), National Institutes for
Quantum and Radiological Science and Technology (QST)
Japan Atomic Energy Agency (JAEA)
The Graduate School for the Creation of New Photonics Industries (GPI)
The Laser Society of Japan (LSJ)
The Japan Society of Plasma Science and Nuclear Fusion Research (JSPF)
The Physical Society of Japan (JPS)
The Ministry of Education, Culture Sports, Science and Technology Japan (MEXT)

***In Situ* Characterization of XFEL Beam Intensity Distribution and Focusability by High Resolution LiF Crystal X-Ray Detector**

T. Pikuz^{1,4}, A. Faenov^{2,3}, T. Matsuoka³, B. Albertazzi⁴, N. Ozaki^{1,4}, N. Hartely⁴, O. Muray Ricardo Arturo^{1,4}, T. Yabuuchi⁶, H. Habara⁴, S. Matsuyama⁴, K. Yamauchi⁴, Y. Inubushi⁵, T. Togashi⁵, H. Yumoto⁵, Y. Tange⁵, K. Tono⁵, Y. Sato⁴, M. Yabashi^{5,6}, M. Nishikino⁷, T. Kawachi⁷, A. Mitrofanov², D. Bleiner⁸, A. Grum-Grzhimailo⁹, N.N. Rosanov¹⁰, N.V. Vysotina¹⁰, M. Harmand¹¹, M. Koenig^{4,12}, K. A. Tanaka^{1,4}, T. Ishikawa^{5,6}, R. Kodama^{1,3,4}

¹Photon Pioneers Center, Osaka University, Suita, Osaka 565-0871, Japan

²Joint Institute for High Temperatures, Russian Academy of Sciences, Moscow, Russia

³Institute for Academic Initiatives, Osaka University, Suita, Osaka, 565-0871, Japan

⁴Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

⁵JASRI/SPRING-8, Sayo, Hyogo 679-5198, Japan

⁶RIKEN Harima Institute, Sayo, Hyogo 679-5148, Japan

⁷Quantum Beam Science Center, Japan Atomic Energy Agency, Kizugawa, Kyoto 619-0215, Japan

⁸EMPA Materials Science and Technology, St Gallen, CH-9014, Switzerland

⁹Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University Moscow, Russia

¹⁰Vavilov State Optical Institute, Kadetskaya Liniya 5/2, 199053 Saint-Petersburg, Russia

¹¹IMPMC, Université Pierre et Marie Curie, CNRS, 75005 Paris, France

¹²LULI, École Polytechnique, CNRS, CEA, UPMC, 91128 Palaiseau, France

Email: tatiana.pikuz@eie.eng.osaka-u.ac.jp

Abstracts: We present here a new X-rays diagnostic based on using LiF X-ray crystal detectors that is able to perform *in situ* measurements the intensity distribution of X-rays beams with diameters ranging from some microns up to some centimeters with high spatial resolution ($\sim 1\mu\text{m}$). We demonstrated also, that by means of the LiF crystal submicron spatial resolution X-ray detector, *in situ* 3D visualization of SACLA XFEL focused beam profile along propagation, including propagation inside photoluminescence solid materials

At coherent X-ray radiation facilities there is an urgent need for simple and efficient methods for *in situ* beamline metrology. Additionally, measurements of energy distribution of XFEL beams in the caustic of focusing system are very important both for correct evaluation of X-ray fluence in the different cross-sections of such beams and for future improving different focusing systems, which are now typically applied for such purposes. We present here a new X-rays diagnostic based on using LiF X-ray crystal detectors that is able to perform *in situ* measurements the intensity distribution of X-rays beams with diameters ranging from some microns up to some centimeters with high spatial resolution ($\sim 1\mu\text{m}$). This diagnostic have an extremely limited cost and is relatively easy to set up. A first observation using this diagnostic at EH5 in SACLA shows that the presence of a slit in the XFEL beam highly structured it by generating diffraction patterns (Fig.1). An optimization of the distance between the slit and the sample position is presented to minimize the effect of the diffraction pattern induced by the slit. We demonstrated also, that by means of the LiF crystal submicron spatial resolution X-ray detector, *in situ* 3D visualization of SACLA XFEL focused beam profile along propagation, including propagation inside photoluminescence solid materials (See Fig.2), is possible [1].

[1] Pikuz, T. *et al.* Sci. Rep. 5, 17713 (2015).