

## Contribution to the flora of Asian and European countries: new national and regional vascular plant records, 5

Marcin Nobis, Arkadiusz Nowak, Renata Piwowarczyk, Aleksandr L. Ebel, Gergely Király, Maria Kushunina, Alexander P. Sukhorukov, Olga D. Chernova, Laura M. Kipriyanova, Beata Paszko, Alexey P. Seregin, Joanna Zalewska-Gałosz, Magdalena Denysenko, Paweł Nejfeld, Adam Stebel & Polina D. Gudkova

**To cite this article:** Marcin Nobis, Arkadiusz Nowak, Renata Piwowarczyk, Aleksandr L. Ebel, Gergely Király, Maria Kushunina, Alexander P. Sukhorukov, Olga D. Chernova, Laura M. Kipriyanova, Beata Paszko, Alexey P. Seregin, Joanna Zalewska-Gałosz, Magdalena Denysenko, Paweł Nejfeld, Adam Stebel & Polina D. Gudkova (2016): Contribution to the flora of Asian and European countries: new national and regional vascular plant records, 5, Botany Letters, DOI: [10.1080/23818107.2016.1165145](https://doi.org/10.1080/23818107.2016.1165145)

**To link to this article:** <http://dx.doi.org/10.1080/23818107.2016.1165145>



Published online: 12 Apr 2016.



Submit your article to this journal [↗](#)



Article views: 35



View related articles [↗](#)



View Crossmark data [↗](#)



## Contribution to the flora of Asian and European countries: new national and regional vascular plant records, 5

Marcin Nobis<sup>a</sup>, Arkadiusz Nowak<sup>b,c</sup>, Renata Piwowarczyk<sup>d</sup>, Aleksandr L. Ebel<sup>e</sup>, Gergely Király<sup>f</sup>, Maria Kushunina<sup>g</sup>, Alexander P. Sukhorukov<sup>h</sup>, Olga D. Chernova<sup>i</sup>, Laura M. Kipriyanova<sup>j</sup>, Beata Paszko<sup>k</sup>, Alexey P. Seregin<sup>l</sup>, Joanna Zalewska-Gałosz<sup>a</sup>, Magdalena Denysenko<sup>m</sup>, Paweł Nejfeld<sup>n</sup>, Adam Stebel<sup>o</sup> and Polina D. Gudkova<sup>e</sup>

<sup>a</sup>Department of Plant Taxonomy, Phytogeography and Herbarium, Institute of Botany, Jagiellonian University, Kraków, Poland; <sup>b</sup>Laboratory of Geobotany and Plant Conservation, Department of Biosystematics, Opole University, Opole, Poland; <sup>c</sup>Department of Biology and Ecology, University of Ostrava, Ostrava, Czech Republic; <sup>d</sup>Department of Botany, Institute of Biology, Jan Kochanowski University, Kielce, Poland; <sup>e</sup>Department of Botany, Institute of Biology, Tomsk State University, Tomsk, Russia; <sup>f</sup>Institute of Silviculture and Forest Protection, Faculty of Forestry of the University of West Hungary, Sopron, Hungary; <sup>g</sup>Department of Plant Physiology, Biological Faculty, Lomonosov Moscow State University, Moscow, Russia; <sup>h</sup>Department of Higher Plants, Biological Faculty, Lomonosov Moscow State University, Moscow, Russia; <sup>i</sup>Siberian Botanical Garden of Tomsk State University, Tomsk, Russia; <sup>j</sup>Novosibirsk Branch of Institute for Water and Environmental Problems, Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia; <sup>k</sup>Department of Vascular Plant Systematics and Phytogeography, W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków, Poland; <sup>l</sup>Department of Geobotany, Faculty of Biology, M.V. Lomonosov Moscow State University, Moscow, Russia; <sup>m</sup>Department of Plant Cytology and Embryology, Institute of Botany, Jagiellonian University, Kraków, Poland; <sup>n</sup>"Dendrus" Environmental Laboratory, Żywiec, Poland; <sup>o</sup>Department of Pharmaceutical Botany, Medical University of Silesia in Katowice, Sosnowiec, Poland

### ABSTRACT

The paper presents new records for 19 vascular plant species from 14 Eurasian countries. Two taxa (*Siphonostegia chinensis* and *Utricularia macrorhiza*) are reported from Russia, two (*Achnatherum botschantzevii* and *Stipa zalesskii*) from Kyrgyzstan, one (*Allium petraeum*) from Uzbekistan, three (*Crambe orientalis*, *Eleocharis mamillata* and *Geranium pratense* L. *fater pratense. sergievskajae*) from Kazakhstan, two (*Atriplex crassifolia* and *Petrosimonia brachyphylla*) from China, one (*Crambe orientalis*) from Tajikistan, one (*Stipa krylovii*) from India, one (*Agrostis lazica*) from Iraq, two (*Orobancha coerulescens* and *Orobancha zaijaci*) from Armenia, one (*Phelipanche lavandulacea*) from Montenegro, one (*Panicum riparium*) from Bosnia and Herzegovina, Romania and Sweden, one (*Sporobolus vaginiflorus*) from Bosnia and Herzegovina and two (*Ranunculus penicillatus* subsp. *pseudofluitans* and *Scutellaria minor*) from Poland. Three of the taxa presented (*Crambe orientalis*, *Panicum riparium* and *Sporobolus vaginiflorus*) are regarded as alien to the studied areas, whereas the remaining 16 are native elements to the flora of the countries. For each species, synonyms, general distribution, habitat preferences, taxonomy with remarks on recognition and differentiation of the species from the most similar taxa occurring in a given country, as well as a list of recorded localities (often far from the previously known areas) are presented.

### ARTICLE HISTORY

Received 30 January 2016  
Accepted 26 February 2016

### KEYWORDS

Alien species; Asia; chorology; Europe; native species; taxonomy

## Introduction

This paper is the continuation of previous works dedicated to new national and regional vascular plant records (Nobis, Nowak et al. 2014; Nobis, Ebel et al. 2014; Nobis, Nowak et al. 2015; Nobis, Ebel et al. 2015). During field exploration across the vast area of 10 European and Asian countries as well as during taxonomic revisions of herbaria materials of different groups of vascular plants, the authors found some species that are new to the floras of particular countries or to significant regions (provinces or republics). The aim of this paper is to report new records for 19 vascular plant species from 14 Eurasian countries, namely Armenia, Bosnia and Herzegovina, China, India, Iraq, Kazakhstan,

Kyrgyzstan, Montenegro, Poland, Romania, Russia, Sweden, Tajikistan and Uzbekistan. Three of the taxa presented here are regarded as alien to the studied areas, whereas the other 16 are native elements to the flora of the countries.

## New records for Asian countries

### *Achnatherum botschantzevii* Tzvelev (Poaceae)

*Contributors* – Marcin Nobis, Arkadiusz Nowak

### *Distribution and habitat*

*Achnatherum botschantzevii* is an endemic species known to date only from locus classicus, which is

located between the Isfara and Sokh River valleys, in the northwestern part of the Alai Mountains in western Kyrgyzstan (Tzvelev 1974), and from one holotypic specimen. During revision of herbarium materials from the genus *Achnatherum*, we found specimens of *Achnatherum botschantzevii* collected from calcareous rocks near Kok-kul Lake, in the border zone of Kyrgyzstan and Uzbekistan. Individuals of the species can probably be found in both of these countries.

### Taxonomic notes

*Achnatherum botschantzevii* is similar to *Achnatherum turkomanicum* (Roshev.) Tzvelev, however it differs from *Achnatherum turkomanicum* by having shorter culms (15–35 versus 40–120 cm long), shorter lemmas (3–4.5 versus 4–5 mm long) and longer awns (20–25 versus 10–15 mm long). *Achnatherum botschantzevii* is also slightly similar to *Achnatherum longiaristatum* (Boiss. & Hausskn.) Nevski (syn. *Stipa kurdistanica* Bor) from Kurdistan, from which it differs by shorter antheria (3–4.5 versus 5.3–6.2 mm long), bearded versus unbearded anthers and also somewhat longer awns (20–25 versus 15–22 mm long; Freitag 1985).

It is worth noting that in the description of *Achnatherum botschantzevii* (Tzvelev 1974) there is a mistake regarding the length of the lemma. Biometrical studies of both the holotype of *Achnatherum botschantzevii* and the specimens from the new locality, show that they have lemmas 3–4.5 mm long, not 1.8–2.2 mm long (as was given by Tzvelev 1974, 1976). This is important information because, based on the protologue of this taxon (Tzvelev 1974) as well as on the later keys to identification of *Achnatherum* (Tzvelev 1976), correct determination of the taxon is impossible. What is more, specimens having long awns and lemmas longer than 2.2 mm could be described as a new taxon.

### Examined specimens (new record)

KYRGYZSTAN/UZBEKISTAN: UdSSR: Fergana / Alai=Gebirge: Umgebung des Sees Kok-kul c. 9 km südöstlich Schachimardan (Chamsabad); Wiederasen; c. 1800 m s.m., 5 July 1989, K.-F. Günther, H.-J. Zündorf, M. Schnittler 758 (JE) – two sheets.

### *Agrostis lazica* Balansa (Poaceae)

Contributor – Beata Paszko

### Distribution and habitat

*Agrostis lazica* was collected and described by Balansa (1874, 12) based on a gathering from Rize district in northeastern Turkey. Until now, *Agrostis lazica* was recorded from Armenia, Azerbaijan, Georgia, Russia (Dagestan) and Turkey (Bitlis, Kayseri, Rize) (Rzazade 1950; Tzvelev 1976, 2006; Doğan 1985; Nersesian 2004; Kurchenko 2010; Nakhutsrishvili 2013). Gillet 9731 is the first record of *Agrostis lazica* for the Kurdistan

Region of Iraq, and extends its known distribution southwards by about 300 km from the closest locality where it was previously recorded at *Süphan Dağı* (Bitlis, Turkey) (Doğan 1985).

*Agrostis lazica* grows in alpine meadows at about 2500–3300 m above sea level (a.s.l.) (Tzvelev 1976; Doğan 1985; Nakhutsrishvili 2013). According to Zazanashvili, Gagnidze, and Nakhutsrishvili (2000) *Agrostis lazica* is a characteristic element of alpine grasslands in the lower alpine belt, from 2400 to 2750 m. In Iraq it was collected at about 3200 m in a damp alpine meadow.

### Taxonomic notes

The genus *Agrostis* L. in Iraq is represented by three species, *Agrostis gigantea* Roth, *Agrostis stolonifera* L. and *Agrostis olympica* (Boiss.) Bor (Bor 1968). An on-going revision of the *Agrostis olympica* complex and its close relatives in the Himalayan region, Near East and Central Asia (Paszko and Pendry 2013; Paszko 2014a, 2014b; Paszko, unpubl. data) revealed a first record of *Agrostis lazica* Balansa (Gillet 9731) housed at K herbarium, from Ser Kurawa (northeast of Erbil) within Rowanduz district (MRO) in the Kurdistan Region (northeast Iraq). Previously, the Iraqi specimen was misidentified by N.L. Bor as *Agrostis olympica*, and as such was cited by him in *Flora of Iraq* (Bor 1968) and in *Flora Iranica* (Bor 1970).

*Agrostis lazica* is often confused with other species, particularly *Agrostis balansae* (Boiss.) Tzvelev (Doğan 1985). It can also be confused with *Agrostis olympica* (Paszko, unpubl. data), as it was by Bor. *Agrostis lazica* differs from *Agrostis balansae* by its relatively dense panicle, short panicle branches (0.5–1.5 cm) and shorter anthers (0.6–0.8 mm) (Doğan 1985). *Agrostis lazica* differs from *Agrostis olympica* by longer callus hairs and anthers, shorter paleas, usually glabrous lemmas (hairy all over in *Agrostis olympica*), and absence of mucros at the tips of lemmas (2–4 mucros up to 0.5 mm long in *Agrostis olympica*) (Doğan 1985; Paszko, unpubl. data).

### Examined specimens (new record)

IRAQ: [Northeast Iraq, Kurdistan Region, Rowanduz distr. (MRO)], Erbil liwa, Ser Kurawa, damp alpine grassland, altitude (alt.) 3200 m, 11 August 1947, J.B. Gillett 9731 (K000482663).

### *Allium petraeum* Kar. & Kir. (Amaryllidaceae)

Contributor – Alexey P. Seregin

### Distribution and habitat

*Allium petraeum* is a widely distributed species of the Central Asian mountains, where it usually occurs on rocks and stony slopes. The most recent revision by Seregin, Anačkov, and Friesen (2015) reports the species from Kazakhstan, Kyrgyzstan, and China.

In Kazakhstan, where *locus classicus* is situated (Karelin and Kirilow 1842), it is known to occur in Tarbagatai,

Dzungarian Alatau, Trans-Ili Alatau, Chu-Ili Mountains and Karatau (Pavlov and Poljakov 1958). Later, Vvedensky (1971) have identified Karatau plants of *Allium petraeum* as *Allium talassicum* Regel. Multiple records either from Tarbagatai or Sauro-Monrak (Pavlov and Poljakov 1958; Bajtenov and Kamenekaja 1990; Kotuhov, Danilova, and Anufrieva 2011) are erroneous and clearly refer to the resurrected *Allium dshungaricum* Vved. (Seregin, Anačkov, and Friesen 2015). Thereby, correct distribution of the species in Kazakhstan covers Dzungarian Alatau, Trans-Ili Alatau and Chu-Ili Mountains.

In Kyrgyzstan, the species was initially recorded only from Kyrgyz Alatau (Nikitina and Kaschenko 1951) and claimed to be of 'rare occurrence' in the republic. Lazkov and Sultanova (2011) have also reported *Allium petraeum* from Northern Kyrgyzstan.

In China, *Allium petraeum* is known from northwest Xinjiang only (Xu and Kamelin 2000). Egorova (1977) has cited correctly specimens from Dzungarian Alatau and E. Tian-Shan. Her record for Dsharkent Depression based on Regel's collection from Suydun [=Shuiding] is erroneous because the locality is situated in the Ili River valley. Another Chinese record by Egorova (1977) based on Potanin's collection from the Kandagatai River in Paraaltai is phytogeographically problematic and most probably refers to another species.

There were no records of *Allium petraeum* from Uzbekistan (Vvedensky 1941, 1971).

### Taxonomic notes

Three collections from Uzbekistan cited below were incorrectly identified as dwarf *Allium talassicum* Regel. *Allium petraeum* is a much taller plant (30–50 cm) with exserted stamens, long valves, white papery bulb tunics and terete sheaths with prominent ribs.

*Allium petraeum* is a taxonomically heterogeneous species from the section *Oreiprason*, which deserves further studies. Preliminary molecular (internal transcribed spacers) results by Seregin and Friesen (2015) suggest that at least three distinct genetic entities co-exist within this species. The sampled specimens examined showed that the situation might be even more complicated. Egorova (1977) correctly reported the presence of red-flowered plants within *Allium petraeum* s.l. Such plants from Dzungarian Alatau are forming a well-supported clade in molecular analysis (Seregin and Friesen 2015) but they were also collected in Sidzhak (Uzbekistan). Specimens from Chimgan are yellow-flowered and look similar to those from Central Tian Shan in Kyrgyzstan.

Vvedensky (1941) considered 65 wild species for Uzbekistan whereas Khassanov (1992) has listed 103 species. Multiple new records as well as new protologues were published afterwards, but no modern estimates on a number of *Allium* species of the Uzbek flora were published afterwards.

### Examined specimens (new records)

UZBEKISTAN: the Pskem River basin, Sidzhak, 14–15 July 1957, *Pyatayeva* (LE); eastern spur of the Greater Chimgan, s.d., *anonymous* (LE); [cultivated in Gatersleben in Germany, Nr. TAX 5883], origin: Uzbekistan, stream terrace covered with grasses and shrubs in Aksaj on northern slope of Chimgan Massif, locally common (May 2001, leg. R. Fritsch & M. Hoffmann 1755), 11 July 2005, *anonymous* (GAT 20125, barcode GAT2554332).

### *Atriplex crassifolia* C.A. Mey. (Chenopodiaceae–Amaranthaceae)

Contributors – Alexander P. Sukhorukov, Maria Kushunina

### Distribution and habitat

*Atriplex crassifolia* is one of the common plants on the saline substrates ('solonchaks') in semi-deserts of central and eastern Kazakhstan and southern Siberia (Novosibirsk, Omsk provinces and Altai Krai) at the altitudes 0–1000 m a.s.l. The eastern range border was known near Lake Zaysan (Kazakhstan), close to the Xinjiang province in China.

A new record for the native flora of China, not reported by Zhu, Mosyakin, and Clemants (2003), is here reported after the revision of herbarium specimens at PE.

### Taxonomic notes

*Atriplex crassifolia* is a member of *Atriplex* sect. *Teutliopsis*, the group of annual representatives with a  $C_3$ -photosynthetic pathway, isolateral leaf anatomy, mixed male and female flowers arranged in clusters, and herbaceous bract-like covers enclosing female flowers (Moser 1934; Kadereit et al. 2010). *Atriplex crassifolia* is clearly distinguished from morphologically similar species by its rhombic and succulent leaves. Previously, its range was considered to include the Lower Volga region of European Russia (Ivanov 1989; Medvedeva 1996), Central Asia and Northern India (Hooker 1890; Paul 2012). However, the revision of the material in many Asian and European herbaria has shown that all records from European Russia belong to *Atriplex patens* (Litv.) Iljin, and those from Northern Himalaya belong to several  $C_4$ -*Atriplex* species (Sukhorukov 2006), especially *Atriplex tatarica* L. and *Atriplex pallida* (Moq.) Sukhor., formerly known as *Atriplex schugnanica* Iljin (Sukhorukov and Kushunina 2015).

### Examined specimens (new record)

CHINA: Xinjiang, [Ili-Kazakh Autonomous prefecture] near Shiqihu, 1000 m, 9.IX.1931, T.N. Niu 3717 (PE). Original text in Chinese.



***Crambe orientalis* L. (Brassicaceae)***Crambe amabilis* Butkov & Majlun

Contributors – Marcin Nobis, Arkadiusz Nowak, Aleksandr L. Ebel

**Distribution and habitat**

*Crambe orientalis* is native to western Asia, ranging from Turkey to Turkmenistan (Hedge 1968; Nikitin and Geldykhonov 1988). The first record of the alien *Crambe orientalis* in western Tian-Shan is dated from 1922 when a specimen was collected from the foothills between Shymkent Town (Kazakhstan) and Angren River (Uzbekistan). Recently, it was found as spreading in Kazakhstan and Kyrgyzstan (Lazkov and Redina 2007; Lazkov and Sennikov 2014), and further localities of the species were found by us during the exploration in Kazakhstan (see below). The species occurred abundantly on grain fields, abandoned lands, along roadsides and on railway embankments.

*Crambe orientalis* was found in Tajikistan for the first time by us in 2009 and the species was observed at the locality also in 2011, 2013 and 2015. It grows fairly abundantly, spreading along roadsides, on arable fields and especially on fallows in the Mogendarya River valley, near the Mogien and Gezani-Bolo villages. It is a newly established and potentially invasive alien species in Tajikistan.

**Taxonomic notes**

There are three species of *Crambe* known to date from Tajikistan: *Crambe schugnana* Korsh., *Crambe kotschyana* Boiss. and *Crambe gordjaginii* Spryg. & M. Pop. (Yunusov 1978). In accordance with the identification key (Yunusov 1978) of Tajik *Crambe* spp., *Crambe orientalis* can be misidentified with the last two species. Similarly to *Crambe kotschiana* it has white petals; however, the most conspicuous character distinguishing both taxa is the shape of the leaves, which are dentate and deeply pinnately lobed in *Crambe orientalis* and broadly cordate at base, ovate to broadly lobate and dentate in *Crambe kotschiana*. The examined species differs from *Crambe gordjaginii* in having white, not yellow petals.

**Examined specimens (new records)**

TAJIKISTAN: Western Zerafshan Mountains, Mogendarya River valley, near Gezani-Bolo settlement, fallows, roadsides, 29 May 2015, M. Nobis, A. Nowak (KRA)

KAZAKHSTAN: Almaty, near Baum's grove, on gravelly embankment of the railway along the avenue Suyunbay, 43.3224303 N/76.958456 E, 5 July 2014, A. Ebel (TK); c. 290 km northwest of Almaty, roadside, near southwest part of the Balkhash Lake, 21 May 2014, observation M. Nobis.

***Eleocharis mamillata* (H. Lindb.) H. Lindb. (Cyperaceae)**

Contributor – Aleksandr L. Ebel

**Distribution and habitat**

*Eleocharis mamillata* s.str. is a holarctic species quite common in northern, central and eastern Europe (Walters 1980; Gregor 2003; Egorova 2007). In Asia, this species is diffusely distributed mainly in Siberia, predominantly south of 60° N, and in adjacent areas of Russian Far East and northern Mongolia (Bubnova 1990; Gubanov 1996; Egorova 2001; Ebel 2012). Recently, it has been discovered in North America where it seemed to be previously confused with *Eleocharis palustris* (L.) Roem. & Schultes and *Eleocharis macrostachya* Britton (Smith et al. 2002). It usually grows in shallow waters, swampy watersides, and on grassy swamps.

*Eleocharis mamillata* is a rare native species in Kazakhstan. It was not listed in the flora of Kazakhstan to date (Poljakov 1958; Abdulina 1999; Egorova 2001) although one location in Central Kazakhstan based on a specimen collected in 1929 in Ulutau by N. Pavlov was mentioned at least twice (Alekseev 1991; Gregor 2003).

**Taxonomic notes**

Limits of *Eleocharis mamillata* are debatable. *Eleocharis mamillata* s.l. comprises two or three subspecies: subsp. *mamillata*, subsp. *austriaca* (Hayek) Strandh. (= *Eleocharis austriaca* Hayek), and East Asiatic subsp. *ussuriensis* (Zinserl.) T.V. Egorova (= *Eleocharis ussuriensis* Zinserl., *Eleocharis austriaca* subsp. *ussuriensis* (Zinserl.) T.V. Egorova). The last one is not accepted by some authors who consider it a synonym of *Eleocharis mamillata* (e.g. Gregor 2003) or, on the contrary, count it as separate species – *Eleocharis ussuriensis* (The Plant List 2013). We prefer to follow the latest opinion of Egorova (2001, 2007) who recognized *Eleocharis austriaca* (including ssp. *ussuriensis*) and *Eleocharis mamillata* as two different species.

*Eleocharis mamillata* differs from *Eleocharis palustris* in having fragile stems with an orbicular rather than oval cross-section, fewer vascular bundles, dense conical flowering heads, and the number of perianth bristles at the base of the fruit usually exceeding four (e.g. Walters 1980; Egorova 2001).

All cited specimens from Kazakhstan belong to *Eleocharis mamillata* subsp. *mamillata* (*Eleocharis mamillata* s.str.).

**Examined specimens (new records)**

KAZAKHSTAN: Akmolinsk Region, Kokchetaw district, Borovoe, bog with bushes, 4 July 1909, V.F. Semenov (TK); North Kazakhstan Region, Sergeevskij district, village Beloglinka surroundings, lake Malyj Tarangul, southwest shore, 4 July 1983, B.F. Sviridenko (TK).

***Geranium pratense* L. subsp. *sergievskajae* Peschkova (Geraniaceae)**

Contributor – Aleksandr L. Ebel

### Distribution and habitat

*Geranium pratense* subsp. *sergievskajae* is an Asian subspecies, partly substituting the Eurasian subspecies *Geranium pratense* subsp. *pratense* in continental regions of Asian Russia. It is widely distributed in Siberia, from Russian Altai and south of Tomsk Region in western Siberia to Buryatia and Sakha (Yakutia) in eastern Siberia (Peschkova 1996; Baikov 2005, 2012). Although this taxon was recognized at the level of subspecies almost 20 years ago, no location in adjacent countries beyond Russia has been recorded so far (Peschkova 1996; Abdulina 1999; Urgamal et al. 2014). It grows predominantly in mountain areas, in coniferous and deciduous forests (dominated by *Pinus sylvestris* L., *Larix* spp. and *Betula* spp.) frequently on forest margins, in river valleys, sometimes in upland meadows, usually in conditions of high humidity of soils and atmosphere.

*Geranium pratense* subsp. *sergievskajae* is a new native taxon to Kazakhstan. Specimens of this subspecies were revealed among materials of *Geranium pratense* collected in the first half of the twentieth century. Moreover, at TK there is an old specimen from Omsk Region (Russia) where this subspecies was not previously recorded.

### Taxonomic notes

This taxon was first separated in 1934 by Sergievskaja as *Geranium pratense* var. *typicum* Kryl. f. *molle* Serg., and much later its status was raised up to subspecies (Peschkova 1996). Only recently was the lectotype of *Geranium pratense* f. *molle* deposited in the Krylov Herbarium (TK) was designated (Troshkina 2015). *Geranium pratense* subsp. *sergievskajae* differs from *Geranium pratense* subsp. *pratense* mainly by type of pubescence. Stems of subsp. *sergievskajae* almost from base are covered by long simple and glandular trichomes more or less of equal length, whereas stems of subsp. *pratense* are covered up to inflorescence by only long simple trichomes. Additionally, leaf blades of subsp. *sergievskajae* are usually more deeply divided into narrower lobes than subsp. *pratense*. The two subspecies also differ in the sculpture of exine and ornamentation of the pollen grain surface (Ivleva 2010).

### Examined specimens (new records)

KAZAKHSTAN: Akmolinsk Region, Borovoe, wet boggy meadow with bushes, 26 June 1908, V.F. Semenov (TK); South Altai, Katon-Karagai neighbourhood, canyon Sarymsak, meadows along riverside, 27 June 1928, V. Evseenko (TK).

### *Orobanche coerulescens* Stephan in Willd. (Orobanchaceae)

Contributors – Renata Piowarczyk, Magdalena Denysenko

### Distribution and habitat

*Orobanche coerulescens* is a Eurasian holoparasite species, occurring from eastern Germany, through eastern Europe and from central Asia to Japan (Pusch 2009; Piowarczyk & Przemyski 2009). It parasitizes *Artemisia* spp., mainly in the sandy or rocky grasslands or *Artemisia* steppes (Piowarczyk 2012). From the Caucasus, the species was known only from Dagestan (Novopokrovskij and Tzvelev 1958) and recently a new locality has been found in the Lesser Caucasus in Georgia (Piowarczyk, and Kwolek in Nobis, Ebel, et al. 2015). New localities of *Orobanche coerulescens* have been found in the Lesser Caucasus in Armenia, and it is a new, native taxon to the flora of this country.

### Taxonomic notes

The flora of Armenia consists of 39 species of *Orobanche* s.l. (Takhtajan 1987). The distribution of this genus and its hosts in Armenia and in the whole area of the Caucasus is not sufficiently known, as shown by new findings (e.g. Rätzel and Uhlich 2004; Piowarczyk 2015a; Piowarczyk et al. 2015).

*Orobanche coerulescens* was included in the section *Inflatae* (Beck) Tzvelev, subsection *Coerulescentes* (Beck) Teryokhin (Teryokhin et al. 1993). Taxonomic problems and relations with similar species were described in detail in previous works (Piowarczyk 2015b; Piowarczyk and Kwolek in Nobis, Ebel, et al. 2015; Piowarczyk, Madeja, and Nobis 2015).

### Examined specimens (new record)

ARMENIA: Northern Armenia, Lesser Caucasus, Gegharkunik province, between Semyonovka and Tsovagyugh, rocky grasslands, on slope with southwestern exposure, parasitizes *Artemisia* spp., alt. 2005 m, 9 July 2015, R. Piowarczyk (KTC).

### *Orobanche zajaciorum* Piow. (Orobanchaceae)

Contributor – Renata Piowarczyk

### Distribution and habitat

*Orobanche zajaciorum* was recently described as a probably endemic species to the Caucasus (Piowarczyk 2015a). It was known only from one locality in the Lesser Caucasus in Georgia. It grows on rocky grasslands and parasitizes *Scutellaria sosnowskyi* Takht. (Piowarczyk 2015a). It is not excluded that the species may parasitize known hybrids of *Scutellaria sosnowskyi* (Çiçek and Yaprak 2011), or other species of the genus *Scutellaria* L., and family Lamiaceae. The new locality of *Orobanche zajaciorum* has been found in the Lesser Caucasus in Armenia, 180 km southeast of the *locus classicus*. The species is a new, native taxon to the flora of this country. The population of the species is small, comprising only five shoots, and is located outside protected areas.

### Taxonomic notes

*Orobanche zajaciorum* has been included in the section *Orobanche* Teryokhin subsection *Galeatae* (Beck) Teryokhin. The corolla shape of *Orobanche zajaciorum*, especially due to the helmet-shaped upper lip, is very similar to that of *Orobanche caryophyllacea* Sm., *Orobanche teucarii* Holandre or *Orobanche lutea* Baumg. However, these three taxa are much taller (usually up to 20 cm) and bear larger flowers (usually 18–30 mm long), bidentate and broad calyx segments, and parasitize different hosts, while *Orobanche zajaciorum*, is usually 10–17 cm high, with distinctively curved geniculate and small flowers (up to 17 mm, usually < 15 mm), simple and narrow calyx segments. These characters make it difficult to confuse it with other taxa of that subsection (Piowarczyk 2015a).

### Examined specimens (new record)

ARMENIA: Northern Armenia, Lesser Caucasus, Gegharkunik province, northwest of Lake Sevan, between Semyonovka and Tsovagyugh, rocky grasslands, on slope with southwestern exposure, parasitize *Scutellaria* cf. *sosnowskyi*, alt. 2005 m, 9 July 2015, R. Piowarczyk (KTC).

### *Petrosimonia brachyphylla* (Bunge) Iljin (Amaranthaceae)

Contributors – Alexander P. Sukhorukov, Maria Kushunina

### Distribution and habitat

*Petrosimonia brachyphylla* is widely distributed in Kazakhstan with extensions into the steppes and semi-deserts of European Russia (Sukhorukov 2014). The westernmost recorded specimen was collected by A. Sukhorukov in Kalmykia in 2012 (G, LE, MW). The eastern range border was known near Zaysan and Balkhash lakes (Kazakhstan). *Petrosimonia brachyphylla* prefers clayey saline soils and often grows together with other native *Petrosimonia* species or other annual halophytes (Sukhorukov 2005). A cited record from Xinjiang is the first one for China.

### Taxonomic notes

The identification of *Petrosimonia* representatives is difficult, and herbarium specimens are often misidentified. The most distinctive features of the genus are the indumentum type, the length of lower leaves (however, they are present in young plants only), the bract/perianth length ratio, the shape of the bracteoles, the number of perianth segments and stamens, and the anthers that can be fused or not fused apically. Short (up to 2 cm), slightly pubescent leaves and apically gibbous bracts surrounding the fruit distinguish *Petrosimonia brachyphylla* from closely related *Petrosimonia crassifolia* (Pall.) Bunge and *Petrosimonia glauca* Bunge, which also have

two perianth segments and five stamens with apically fused anthers.

### Examined specimen (new record)

CHINA: Xinjiang, Altay, Salhushun, alt. 450 m, 17 September 2008, Zhai Wei, Wang Guangfu 2405 (SHI).

### *Siphonostegia chinensis* Benth. (Orobanchaceae)

Contributor – Olga D. Chernova

### Distribution and habitat

*Siphonostegia chinensis* is an annual species native to East Asia (Starchenko 2008). It occurs in China, Japan, Korea (Hong et al. 1998) as well as in the far east of Russia. For the Russian territory, it is only reported in Manchurian continental and Sakhalin oceanic floristic provinces (Ivanina 1991; Baikov 2012). The species is not reported for Siberian Flora (Baikov 2005). It grows in steppes, on rocky hills, scrublands, light forests, meadows and sod fields (Ivanina 1991). In China, it is typical of dry mountain slopes and meadows at a height between 800 and 3400 m (Hong et al. 1998).

*Siphonostegia chinensis* is a new native species for Siberia. The species has been found in Trans-Baical Territory (at 3 km and 17 km from the border with China, Inner Mongolia Autonomous Region) on mid-mountain lots of the southeastern spurs of the Gazimur Range, in the lower course of the river Budyumkan (left tributary of the river Argun) within petrophilous Sheep Fescue associations in the upper part of steep steppe slopes. According to the zonation of Asian Russia (Baikov 2012), this territory belongs to the Baikal hemiboreal province.

### Taxonomic notes

The genus *Siphonostegia* Benth. includes two or three annual East Asian species and one eastern Mediterranean perennial species, the latter one sometimes separated into monotypic genus *Lesquereuxia* Boiss. & Reut. (Hong et al. 1998; Fischer 2004). All of them are hemiparasitic plants. *Siphonostegia chinensis* is a single species of the genus having bi-pinnatisect leaves.

### Examined specimens (new records)

RUSSIA: Zabaykalye Territory, Gazimuro-Zavodskoy district, Relic Oaks State Reserve, 6 km south of Uryupino outpost, left bank of the river Budyumkan, upper part of the steppe rubble slope at southwestern exposition, 52°41' N, 120°01' E; alt. 425 m, 19 July 2010, O.D. Chernova (TK); Zabaykalye Territory, Gazimuro-Zavodskoy district, 3 km from Budyumkan village, road fork district, left bank of the river Budyumkan, upper part of the steppe rubble slope at southern exposition, 52°37' N, 119°47' E; alt. 486 m, 21 July 2013, O.D. Chernova (TK).



### *Stipa krylovii* Roshev. (Poaceae)

*Stipa capillata* var. *coronata* Roshev., *Stipa densiflora* P.A.Smirn., *Stipa densa* P.A.Smirn., *Stipa decipiens* P.A.Smirn., *Stipa sareptana* subsp. *krylovii* (Roshev.) D.F.Cui, *Stipa sareptana* var. *krylovii* (Roshev.) P.C.Kuo & Y.H.Sun

Contributors – Marcin Nobis, Arkadiusz Nowak, Polina Gudkova

### Distribution and habitat

*Stipa krylovii* is a widely distributed central Asian taxon, ranging from southern Russia (Siberia) through Kazakhstan, Mongolia, Kyrgyzstan, Tajikistan and China, to Tibet, Nepal and northeastern India (Tzvelev 1976; Wu and Phillips 2006; Gudkova and Nobis unpubl. data). Occurrence of the species in India is given for the first time by us. Specimens of this taxon were found in Ladakh by L. Klimeš, but identified as *Stipa capillata* L. *Stipa krylovii* is a new native taxon to the flora of India. It occurs on high mountain steppes and semi-deserts.

### Taxonomy

To date, c. 20 taxa of Stipaeae are known from India (Bor 1960; Cope 1982; Freitag 1985; Dickoré 1995; Klimeš and Dickoré 2005; Nobis 2011, 2014; Nobis, Nobis, et al. 2014). *Stipa krylovii* is close to *Stipa capillata* L., but differs in having shorter ligules of the vegetative shoots (0.1–0.3 versus 0.6–2.5 mm long, respectively), thinner leaves (0.3–0.5 versus 0.5–1.2 mm wide, respectively) and the presence of a well-developed ring of hairs on the top of lemma (in *Stipa capillata* top of anthercium is glabrous, rarely with single short prickles). *Stipa krylovii* can also be confused with *Stipa sareptana* A.K.Becker. The two taxa differ mainly in the character of vegetative leaves (glabrous to slightly scabrous in *Stipa krylovii* versus distinctly scabrous covered by tubercles and spinules in *Stipa sareptana*), somewhat shorter ligules of the vegetative shoots (0.1–0.3 versus 0.2–1.0 mm long, respectively) as well as in the pattern of general distribution (Tzvelev 1976). There are conflicting opinions about the systematic position of *Stipa krylovii*, but most researchers treat it at the specific level (Roshevitz 1934; Pazij 1968; Tzvelev 1976; Lomonosova 1990; Nobis and Gudkova 2016).

### Examined specimen (new record)

INDIA NW: Jamu and Kashmir State, Ladakh, Indus Vy: Zhung (Leh), Chkirmo, 33°49.5' N/77°39.1 E, alt. 4150 m a.s.l., 8 September 2001, L. Klimeš 1626 (PR).

### *Stipa zalesskii* Wilensky (Poaceae)

Contributors – Marcin Nobis, Arkadiusz Nowak

### Distribution and habitat

*Stipa zalesskii* is a widely distributed taxon ranging from central Europe to central Asia (Kazakhstan, Russia)

(Martinovský 1980; Conert 1998; Tzvelev 1976, 2012; Freitag 1985). In central Asia, the species is known to date only from Kazakhstan. The species is not mentioned in the checklist flora of Kyrgyzstan (Lazkov and Sultanova 2011). During field studies in Kyrgyzstan in 2013 and 2015 we found several abundant populations in the Sasuumyr River valley on the bottom of the Kyrgyzian Mountains and Talas Alatau. This is a new, native species in the flora of the country.

### Taxonomy

In the flora of Kyrgyzstan, there are c. 30 species of *Stipa* (Lazkov and Sultanova 2011; Nobis 2012, 2013; Nobis, Ebel, et al. 2014, 2015; Nobis, et al. 2016). Localities of *Stipa zalesskii* in Kyrgyzstan are situated on the easternmost range limit of the taxon. *Stipa zalesskii* belongs to the section *Stipa* and is well distinguished from others Kyrgyz feather grasses from the section, namely *Stipa kirghisorum* P.A.Smirn., *Stipa macroglossa* P.A.Smirn. subsp. *macroglossa*, *Stipa macroglossa* subsp. *kazakhstanica* (Kotukhov) M.Nobis, *Stipa turkestanica* Hack. subsp. *trichoides* (P.A.Smirn.) Tzvelev (Nobis et al. 2016) by the longest anthercia (18.5–23 cm) and the ventral line of hairs on the anthercium reaching the base of the awn or ending at 0.3 mm below the base of awn.

According to Tzvelev (1976) *Stipa zalesskii* is characterized by having the lemma 16–19 mm long, ventral line of hairs on lemma reaching or at 0.5–1.3 mm not reaching the base of the awn, leaves of the vegetative shoots with mixed short and long hairs at the adaxial surface and scabrous with more or less dense admixture of spinules or short hairs at the abaxial surface. However, the specimens of *S. zalesskii* found by us in Kyrgyzstan have distinctly longer anthercia (18.5–23 mm long), which make them similar to *Stipa turkomanica* P.A.Smirn. [= *Stipa zalesskii* subsp. *turkomanica* (P.A.Smirn.) Tzvelev] occurring in Turkmenistan. The only difference is the length of awn seta, which is generally shorter in *Stipa zalesskii* (up to c. 35 cm versus over 35 cm long respectively) and character of sheaths of the vegetative shoots, which are shortly pubescent in Kyrgyz specimens versus glabrous in *Stipa turkomanica*. This character (pubescence or its lack on sheaths) is often considered as a natural variation, and glabrous and pubescent sheaths can be observed within one population; however, after preliminary revision of the herbarium materials belonging to both of these taxa at LE, MW, KRA, TK and MHA, we did not note such a situation in specimens identified as *Stipa zalesskii* and *Stipa turkomanica*. Nevertheless, some researchers treat *Stipa turkomanica* as conspecific with *Stipa zalesskii* (Freitag 1985). In the *Stipa zalesskii* group belong such morphologically similar taxa as *Stipa dasyphylla* (Lindem.) Trautv., *Stipa rubens* P.A.Smirn., *Stipa rubentifformis* P.A.Smirn., *Stipa glabrata* P.A.Smirn., *Stipa pontica* P.A.Smirn., *Stipa canescens* P.A.Smirn. ex Roshev., *Stipa iljinii* Roshev., *Stipa maeotica* Klokov



and *Ossuczniuk*, and *Stipa smirnovii* Martinovský. This group of taxa requires taxonomic revision (Nobis et al. unpubl. data).

### Examined specimens (new records)

KYRGYZSTAN: steppe grassland, 20 km west-northwest of Suusamyр (Suusamyр River valley, Talasski Alatau), roadside, alt. 2250 m a.s.l., slope 20°, inclination south, 7 July 2015, *M. Nobis*, *A. Nowak* (KRA); steppe grassland, 27 km west-northwest of Suusamyр (Suusamyр River valley, Talasski Alatau), by the road, alt. 2280 m a.s.l., slope 40°, inclination southwest, 30 June 2015, *M. Nobis*, *A. Nowak* (KRA); steppe grassland, 12 km northwest of Suusamyр (escarpement by the road), alt. 2312 m a.s.l., slope 15°, inclination southwest, 7 July 2015, *M. Nobis*, *A. Nowak* (KRA); steppe grassland, 11 km northwest of Suusamyр (by the road), alt. 2287 m a.s.l., slope 5°, inclination northwest, 7 July 2015, *M. Nobis*, *A. Nowak* (KRA); steppe grassland, 7 km northwest of Suusamyр (escarpement by the road), alt. 2010 m a.s.l., slope 25°, inclination east, 7 July 2015, *M. Nobis*, *A. Nowak* (KRA); Talasskii Alatau, grasslands by the main road in the Suusamyр River valley, alt. 2270 m a.s.l., slope 30°, inclination south, 30 June 2013, *M. Nobis* (KRA).

### *Utricularia macrorhiza* Le Conte (Lentibulariaceae)

Contributor – Laura M. Kipriyanova

### Distribution and habitat

*Utricularia macrorhiza* is distributed in North America and temperate regions of Asia: Kamchatka, Sakhalin, Amur and Ussuri regions, Mongolia and China (Taylor 1989). The author noted that the species extends west to the Altay; however, the observation was not based on relevant herbarium samples. Recently, this species was found as quite common in East Siberia (Chepinoga and Rosbakh 2012; Pazdnikova and Chepinoga 2013) as well as in West Siberia in Khanty-Mansi Autonomous District – Yugra (Kapitonova, Kapitonov, and Ilminskikh 2014). The authors suggested that *Utricularia macrorhiza* is likely to be common in aquatic habitats in West Siberia, but that it is not distinguished by collectors due to external similarities with *Utricularia vulgaris* L. (Kapitonova, Kapitonov, and Ilminskikh 2014). *Utricularia macrorhiza* was also not included in the manual of the Siberian species (Olonova 1996) and other regional manuals, and the species was skipped by researchers also on those grounds.

*Utricularia macrorhiza* was confirmed in the three lakes of Altai Region (West Siberia) namely: Lena, Bol'shoe, and Rakity in 2014. The salinity of the water from Lena Lake was 4.2 g/dm<sup>3</sup>. In this lake only *Chara tomentosa* L. was noted in the community with the dominance of *Utricularia macrorhiza*. Salinity of the Bol'shoe Lake was 0.262 g/dm<sup>3</sup>. The lake is transformed into a

swamp, and bladderwort was found in the water among the floating communities of *Thelypteris palustris* Schott, *Carex diandra* Schrank, *Phragmites australis* (Cav.) Trin. ex Steud. *Hydrocharis morsus-ranae* L., *Utricularia minor* L. and other aquatic plants were found with *Utricularia macrorhiza*. Salinity of the Rakity Lake was 0.689 g/dm<sup>3</sup>, the bladderwort formed coenoses in the near-shore zone with *Lemna minor* L. Plants were blooming in all the lakes; some of them were beginning to form fruits.

### Taxonomic notes

*Utricularia macrorhiza* differs from *Utricularia vulgaris* mainly by the distal half of the spur, which is narrow and cylindrical. The spur at the end is relatively acute and visibly bent upward, while the distal half of the *Utricularia vulgaris* spur is shortly conical and straight with a blunt tip (Taylor 1989). It should be noted that it is very useful to take pictures in nature, because the flowers in the herbarium often fall off or become deformed.

There are glands on the inside of the dorsal and ventral surfaces of the spur of *Utricularia macrorhiza* (Lisitsyna and Papchenkov 2000; Taylor 1989) whereas in *Utricularia vulgaris*, the glands develop only on the inside of the dorsal surface. However, the collection should be representative enough because an attribute based on glands is not well expressed in each flower. Some variations in the severity of attributes appear to be associated with an overlapping of the areas of distribution of the two species. Lisitsyna and Papchenkov (2000) note that *Utricularia macrorhiza* is a more 'gentle' plant; the extreme segments of the leaves are subulate-filiform with few spines on the surface and ending in spines whereas in the case of the more rigid *Utricularia vulgaris*, the extreme segments of leaves are covered with sparse cilia.

### Examined specimens (new records)

RUSSIA: Altai Region, Baevsky district, lake Lena, 2 km to the northwest of Baevo village, 53°16'07.5" N, 80°41'00.1" E, 6 July 2014, *L.M. Kipriyanova* (NS); Altai region, Uglovsky district, lake Bol'shoye close to Lyapunovo village, 51°18'30.7" N, 80°09'28.7" E 13 July 2014, *L.M. Kipriyanova* (NS); Altai region, Rubtsovsky district, lake Rakity, 51°33'30.5" N 81°11'49.8" E, 14 July 2014, *L.M. Kipriyanova* (NS).

### New records for European countries

#### *Panicum riparium* H. Scholz (Poaceae)

Contributor – Gergely Király

### Distribution and habitat

In Europe, *Panicum riparium*, an adventive taxon originated from North America, was recognized for the first time in Germany (Scholz 2002); later, it was observed in several other countries: Austria (Hohla 2006), Hungary

(Király et al. 2009), Switzerland (Ciardo et al. 2011), Italy (Wilhelm 2011), France and Great Britain (Amarell 2013), Belgium (Verloove 2014), Croatia (Király and Alegro 2015). However, herbarium revisions showed that this taxon was formerly overlooked, because it has been present – accompanied by *Panicum capillare* – in Europe since the eighteenth century (Amarell 2013; Király and Alegro 2015). *Panicum riparium* grows both in ruderal habitats and (often as a noxious weed) in corn and winter wheat cultures (Clements et al. 2004; Nagy et al. 2012). The species seemed to have a rather sub-Atlantic character and reports of its occurrence in eastern parts of Europe are rare or missing.

In the course of revision of herbarium material of the *Panicum capillare*-complex, specimens of *Panicum riparium* were recognized from Bosnia and Herzegovina (Bihać), and for Romania (Moldova Veche and Gigheria); the species was formerly not recorded in these countries. The Romanian localities represent the easternmost outposts of this species. Amarell (2013) reported a specimen from Sweden (LINN 80.50) but from the botanical garden of Uppsala. Hence, a new herbarium specimen collected in Göteborg in ruderal habitats is the first confirmation of its presence in the wild for Sweden, and, in addition, for Scandinavia.

### Taxonomy

The representatives of the *Panicum capillare* complex (*Panicum* sect. *Panicum*, ‘witchgrasses’) are native to North America; in Europe a single species, *Panicum capillare* L., is considered as an often naturalized alien; other species were recorded as casuals only (Clayton 1980; Ryves et al. 1996; Freckmann and Lelong 2007). A new species of the complex, *Panicum riparium*, was recently described from Germany by Scholz (2002) who emphasized that it has been spontaneously created from American parents. Later, the species was reported under the name *Panicum riparium* from several European countries and seems to be widely accepted. On the other hand, Amarell (2013) argued for *Panicum barbipulvinatum* Nash being an older valid name for this taxon; in his opinion both names are synonyms. Although some authors (e.g. Hohla 2013) have already adopted the name *Panicum barbipulvinatum*, its use is not yet widespread. At present we also prefer to keep the name *Panicum riparium*.

*Panicum riparium* is easily distinguishable from *Panicum capillare* by pedicels of subterminal spikelets shorter than 3 mm and appressed to the branch (*Panicum capillare* has pedicels of subterminal spikelets longer than 5 mm and patent from the branch). The mature spikelets of *Panicum riparium* are 2.7–3.4 times as long as broad, acuminate with a long tapering apex; lemmas are 5(–7) veined. In the case of *Panicum capillare* mature spikelets are 2.2–2.7 times as long as broad, acute with a short pointed apex; lemmas (7–)9

are veined (for further details see Amarell 2013; Király and Alegro 2015).

### Examined specimens (new records)

BOSNIA and HERZEGOVINA: Una-Sana Canton, ‘Wüste Orte bei Bihać’ [Ruderal places near Bihać], n. d., Boller as ‘*Panicum capillare* L., rev. K. Pagitz as ‘*P. barbipulvinatum* Nash’ (WU, Halácsy Herbarium Europaeum). ROMANIA: Caraş-Severin County: ‘Flora Romaniae Exsiccata. In cultis saepe inundatis insulae Ostrov prope Moldova Veche. Alt. cc. 90 m. s. m., 23 September 1968, D. & M. Cîrţu as ‘*Panicum capillare* L., rev. G. Király as ‘*P. riparium* H. Scholz’ (BP 638480 and W 23888). Dolj County: ‘Flora Olteniae Exsiccata. In arenosis ‘*Astragalus contortuplicatus* – *Chlorocyperus glomeratus* ass’ ad ripam fluminis Danubium, inter pagos Gighera et Nedeia. Alt. cc. 15 m. s. m., 1 September 1971, I. Morariu, P. Ularu & M. Danciu as ‘*Panicum capillare* L., rev. K. Pagitz as ‘*P. barbipulvinatum* Nash’ (WU). SWEDEN: ‘Göteborg, Delsjöupplaget, in ruderalis’, 11 October 1953, C. Blom as ‘*Panicum capillare* L., rev. G. Király as ‘*P. riparium* H. Scholz’ (W 13288).

### *Phelipanche lavandulacea* (Rchb.) Pomel (Orobanchaceae)

*Orobanche lavandulacea* Rchb.

Contributor – Renata Piwowarczyk

### Distribution and habitat

*Phelipanche lavandulacea* occurs in the Mediterranean region from the Iberian Peninsula eastwards through southern Europe and northern Africa up to Asia Minor, Syria and Lebanon (Beck von Mannagetta 1930; Carlón, et al. 2015). It is parasitic strictly on *Psoralea bituminosa* L. (Fabaceae). Recently, it was found in Albany (Malo and Shuka 2008), but was probably mistakenly listed on *Thymus*. Some new localities were also found in Croatia (Frajman and Schönschwetter 2008; Piwowarczyk unpubl.). New localities of *Phelipanche lavandulacea* were found in Montenegro, so complementing its range in the southern Balkans. It is a new, native taxon to the flora of this country. In Montenegro, it grows on limestone roadsides, often in ecotone areas between the main roads and rock walls, on rock ledges as well as in ruderal habitats. Populations of the species varied from a few individuals to over 70 shoots.

### Taxonomic notes

*Phelipanche lavandulacea* is easily recognized from the rest of the southeastern European species of *Phelipanche* Pomel, by the often branched and tall stem, deeply dark purple, violet or blue flowers and its specific host. The sub-species *Phelipanche lavandulacea* subsp. *trichocalyx* (Webb & Berthel.) Carlón, G.Gómez, M.Laínz, Moreno Mor., Ó.Sánchez & Schneew. may be only a geographic

race of the Mediterranean *Phelipanche lavandulacea* s. str. in the Canary Islands (cf. Carlón et al. 2008). *Phelipanche lavandulacea* subsp. *trichocalyx* parasitizes also on *Psoralea*, but differs from typical species mainly by calyx teeth, which have the same length as the tube (0.80–1.18), the corolla more intensively violet on the margins of petals, which are also longer, the folds of the corolla and the anthers glabrous (Carlón et al. 2008).

Possible confusion can also arise with the recently described species that are endemic to Spain, from Ebro basin – *Phelipanche lavandulaceoides* Carlón, G. Gómez, M. Laínz, Moreno Mor., Ó. Sánchez & Schneew. (Carlón et al. 2008). Although it shares the same host and some morphological features with *Phelipanche lavandulacea*, it differs substantially both genetically and morphologically (less branched stem, calyx always violet on the entire surface, longer teeth of the calyx and tube 0.94–1.25(–1.5), corolla ± tubular-funnel, clearly widened towards the throat wider, with broadly rhombic-spatulate lobes) (Carlón et al. 2008).

### Examined specimens (new records)

MONTENEGRO: Southern Montenegro, Valdanos Peninsula, near Čivlak, roadsides, edges of scrub, also near the fences of buildings, alt. 80 m, 2 May 2015, R. Piwowarczyk (KTC); between Budva and Boreti, serpentine road towards Cetinje, roadside, ecotone zones of roads and rock walls, steep rock shelves, alt. 197 m, 5 May 2015, R. Piwowarczyk (KTC); Ulcinj, c. 150 m west of the cemetery in the old town, ruderal habitat in the area of ecotone roadside and rocks at buildings, alt. 23 m, 6 May 2015, R. Piwowarczyk (KTC); CROATIA: Southeast Croatia, between Čilipi and Popovići, south of Komaji, rural roadside, alt. 150 m, 8 May 2015, R. Piwowarczyk (KTC).

### *Ranunculus penicillatus* subsp. *pseudofluitans* (Syme) S. D. Webster (Ranunculaceae)

Contributor – Joanna Zalewska-Gałosz

### Distribution and habitat

*Ranunculus penicillatus* subsp. *pseudofluitans* occurs throughout Europe except in the extreme north and the Balkan Peninsula (Cook 1966). It grows in rivers and streams, mainly over calcareous or base-rich substrates (Webster 1988). Although *Ranunculus penicillatus* subsp. *pseudofluitans* is known from Eastern Germany, it has never been recorded in Poland to date. It is a new native taxon for the flora of the country.

### Taxonomic notes

*Ranunculus penicillatus* (Dumort.) Bab. comprises a heterogenic group of aquatic buttercups in the section *Batrachium* (DC) A. Gray, which occur predominantly in swiftly flowing water in rivers and streams (Cook 1966). In terms of formal taxonomy, individual

taxa of *Ranunculus penicillatus* were differently classified. Homophyllous plants, developing long submersed leaves were described as *Ranunculus calcareus* Butcher (1960). The name *Ranunculus pseudofluitans* (Syme) Newbould ex Baker & Foggitt was also applied to such homophyllous plants; however, nowadays, this name is synonymised with *Ranunculus penicillatus* subsp. *penicillatus*. In the monograph of *Batrachium*, Cook (1966) distinguished three varieties within *Ranunculus penicillatus*, namely var. *penicillatus*, var. *calcareus* (Butcher) C.D.K. Cook and var. *vertumnus* C.D.K. Cook. Later, Webster (1988) recognized two subspecies within the group: subsp. *penicillatus* and subsp. *pseudofluitans* (Syme) S.D. Webster. Webster's approach (1988) is now widely accepted and it is also applied here.

*Ranunculus penicillatus* subsp. *pseudofluitans* is most similar to *Ranunculus fluitans* Lam.; however, some differences allow the differentiation of both taxa. The leaves of *Ranunculus penicillatus* subsp. *pseudofluitans* are more divided than those of *Ranunculus fluitans* (often seven or eight divisions versus up to four divisions). Also, *Ranunculus fluitans* rarely forms roots at the nodes whereas subsp. *pseudofluitans* usually does. In the generative stage, differentiation is more straightforward. *Ranunculus fluitans* has receptacles that are totally glabrous or with only a few hairs, whereas those of subsp. *pseudofluitans* are densely pubescent.

*Ranunculus penicillatus* subsp. *pseudofluitans* is morphologically very variable. To reflect this variation, Webster (1988) distinguished two varieties within the subspecies; a robust one, with leaves longer than adjacent internodes – var. *pseudofluitans* (Syme) S.D. Webster and a smaller one, with more divided leaves, conspicuously shorter than adjacent internodes – var. *vertumnus* C.D.K. Cook.

Individuals from Poland are also very variable and a continuous series of intermediates between both morphotypes can be observed. In the river Płytnica, robust plants similar in appearance to *Ranunculus fluitans* were noticed whereas in the river Piława, small morphotypes with dense and relatively short leaves were evidenced. Moreover, in the river Rurzyca, morphologically variable individuals grew side by side. Hence, the great variation suggests a hybridogenous origin for this taxon, which was previously suggested for all taxa from the *Ranunculus penicillatus* group (Cook 1966; Holmes 1979) and which, for some of them, was recently evidenced based on molecular studies (Zalewska-Gałosz, Jopek, and Ilnicki 2015; Bobrow et al. 2015). Probably, *Ranunculus penicillatus* subsp. *pseudofluitans* is also of hybridogenous origin, but this demands further studies.

### Examined specimen (new record)

POLAND: northwestern Poland, river Piława in Szwecja village, 53°21'09" N, 16°34'18.62" E, 13 August 2013, J. Zalewska-Gałosz (KRA); northwestern Poland, river Płytnica in Płytnica village, 53°18'16.4"



N, 16°47'31.45" E, 13 August 2013, J. Zalewska-Gałosz (KRA); northwestern Poland, river Rurzyca in Krępsko village, 53°15'48.3" N, 16°46'38.8" E, 13 August 2013, J. Zalewska-Gałosz (KRA).

### *Scutellaria minor* Huds. (Lamiaceae)

Contributors – Paweł Nejfeld, Adam Stebel

#### Distribution and habitat

*Scutellaria minor* is a species with oceanic distribution (Ellenberg et al. 1992). In Europe, it is found in the Azores, Belgium, Great Britain, France, Germany, Holland, Ireland, Italy, Portugal, Spain, Sweden and Austria [<http://www.ecoflora.co.uk>] (Fitter and Peat 1994). In Poland, *Scutellaria minor* has been classified as a temporarily introduced alien plant species (ephemero-phyte) (Urbisz 2011; Tokarska-Guzik et al. 2012), with a single location near Wolsztyn in the Greater Poland Voivodeship, reported by Szulczewski (1951). Careful literature studies pointed out that the species is rather a native one and was reported only in the nineteenth century, on two sites: the bank of Święte Lake near Wolsztyn in Greater Poland Voivodeship (Hellwig 1897, cited by Szulczewski 1951) and a place called 'Fasanen Garten' between Jeziory Dolne and Suchodół villages, near Lubsko in Lubusz Voivodeship (Baenitz 1861; Decker 1911). Probably, nowadays, both sites do not exist. The wet meadows between Jeziory Dolne and Suchodół were overgrown by tall herb and thicket communities (S. Rosadziński pers. comm.) and in the regional red list of Greater Poland Voivodeship, this species was given the DD category (Jackowiak et al. 2007).

A new, abundant locality (about 250 individuals) of *Scutellaria minor* has been found in Silesia Province, southwestern Poland. *Scutellaria minor* grows there in sedge mire. The floristic composition in the patch of community with *S. minor* has been studied, with relevés using a standard Braun-Blanquet method. Vascular plant names follow Mirek et al. (2002), while mosses follow Ochrya et al. (2003). Area of 25 m<sup>2</sup>, cover of the herb layer c: 90%, cover of the moss layer d: 90%, number of species in the relevé: 32. *Scutellaria minor* 1; *Sphagnum palustre* 4; *Carex echinata* 3; *Potentilla erecta* 3; *Sphagnum flexuosum* 3; *Briza media* 2; *Carex panicea* 2; *Drosera rotundifolia* 2; *Epipactis palustris* 2; *Equisetum fluviatile* 2; *Holcus lanatus* 2; *Sanguisorba officinalis* 2; *Sphagnum teres* 2; *Anthoxanthum odoratum* 1; *Carex nigra* 1; *Cirsium palustre* 1; *Dactylorhiza fuchsii* 1; *D. majalis* 1; *Eriophorum angustifolium* 1; *E. latifolium* 1; *Festuca rubra* 1; *Juncus articulatus* 1; *Lotus uliginosus* 1; *Lysimachia vulgaris* 1; *Nardus stricta* 1; *Crepis paludosa* +; *Danthonia decumbens* +; *Epilobium palustre* +; *Equisetum palustre* +; *Galium uliginosum* +; *Juncus effusus* +; *Myosotis palustris* +.

The floristic composition of this phytocoenosis with *Scutellaria minor* corresponds with the phytosociological

preferences of the species in literature (Ellenberg 1996; Costa et al. 2012). It is most frequently recorded within the plant associations with Atlantic and sub-Atlantic range like *Anagallido tenellae-Juncion bulbosi* alliance from *Scheuchzerio-Caricetea nigrae* and *Juncion acutiflori* alliance from the *Molinio-Arrhenatheretea* class. The newly discovered station in Poland is located at the eastern distribution limit of the species in Europe. In the light of the new data, *Scutellaria minor* seems to be a native species in the flora of Poland.

#### Taxonomic notes

In Poland, the genus *Scutellaria* comprises four species: *Scutellaria altissima* L., *Scutellaria galericulata* L., *Scutellaria hastifolia* L., and *Scutellaria minor*. *Scutellaria minor* differs from the most similar *Scutellaria galericulata* mainly by the colour and shape of corolla, which is 6–7(–10) mm long and pink to pinkish-purple with straight tube in *Scutellaria minor* versus 15–22 mm and blue, violet, rarely pink with folded tube at an angle of about 45° in *Scutellaria galericulata* and leaves predominantly entire or 1–2 pairs slightly serrated in lower part of stem in *Scutellaria minor* versus all leaves slightly serrated in *Scutellaria galericulata*. The hybrids between *Scutellaria minor* and *Scutellaria galericulata*, *Scutellaria* × *hybrida* Strail, are observed.

#### Examined specimens (new record)

POLAND: Western Carpathians, Beskid Mały range, Kocoń village near Ślemień, 49°44'05"N/19°23'58" E, alt. 575 m, exp. south, slope 7°, in the community from the *Scheuchzerio-Caricetea nigrae* class, 22 July 2015, A. Hulbój, P. Nejfeld (KRA).

### *Sporobolus vaginiflorus* (Torr. ex A. Gray) Alph. Wood. (Poaceae)

Contributor – Gergely Király

#### Distribution and habitat

*Sporobolus vaginiflorus* is a grass species native to North America (Hansen 1980; Peterson, Hatch and Weakley 2007), which was first reported from Europe (Italy) in 1951 (Wilhelm 1998). Due to its rapid expansion, it later reached more Southeastern and Central European countries; the furthest points of its European range are now in southern Germany and Hungary (Fürnrohr 2015; Király and Hohla 2015). In some regions of moderate climate, it is classified as an invasive species (Melzer 2003; Tinner 2013).

*Sporobolus vaginiflorus* has not been reported from Bosnia and Herzegovina, but it is probably not rare and is only overlooked (in the neighbouring Croatia it has been known since the 1950s; Horvatić and Gospodarić 1960). On the first recorded locality in the northern part of the country (near Bosanski Petrovac), the species composes homogeneous stands

on the shoulders of an asphalt road at a considerably high elevation (near 800 m a.s.l.). Detailed phytocoenological studies on European *Sporobolus vaginiflorus* stands have not yet been published, but all floristic reports mentioned it from species-poor ruderal places. Conservation threats caused by the species have not yet been reported.

### Taxonomy

The genus *Sporobolus* involves two rather rare species native to the European Mediterranean. Additionally, several exotic representatives of the genus were recognized in Europe; however, only two species: *Sporobolus neglectus* Nash and *Sporobolus vaginiflorus* were reported to be invasive in the southeastern part of the continent (Alegro et al. 2003; Melzer 2003; Hohla, Diewald and Király 2015).

*Sporobolus vaginiflorus* is an unimpressive, morphologically variable species (Harms 2015). It is distinguishable from the only similar representative of the genus (*Sporobolus neglectus*) in Europe by the strigose lemmas, by spikelets 2.5–6.0 mm long, and by mature fruits longer than 1.8 mm. The uppermost leaf sheet overhangs the top of the inflorescence.

### Examined specimen (new record)

BOSNIA and HERZEGOVINA: Una-Sana Canton, 3 km west of Jasenovac along the road to Bosanski Petrovac, roadsides, alt. 785 m, 44°33'29" N, 16°30'14" E, 29 September 2013, G. Király, Z. Barina & D. Pifkó (BP).

### Acknowledgements

Alexey P. Seregin is deeply indebted to Dr Klaus Pistrick, a curator of the GAT herbarium for support and hospitality during the visit. The researches of Alexey P. Seregin, Alexander P. Sukhorukov and Maria Kushunina were supported by Russian Scientific Foundation (RNF, grant #14-50-00029). The research of Joanna Zalewska-Gałosz was funded by the National Science Centre, Poland (DEC-011/01/B/NZ8/00099). Research of Aleksandr L. Ebel was supported by the D.I. Mendeleev Scientific Fund Programme of Tomsk State University. Research of Laura Kipriyanova was supported by the Russian Foundation for Basic Research (grants NN 13-04-02055 a, 14-04-10164-k, 15-29-02498-ofi\_m). The study of Gergely Király was supported by the project 'Agrárklíma.2 VKSZ-12-1-2013-0034'. Research visits of Beata Paszko to herbaria were supported as follows: in 2013 to St Petersburg (LE) by the exchange programme between the Polish Academy of Sciences and the Russian Academy of Sciences, in July 2015 to Edinburgh (E) and London (K) by the SYNTHESYS Project financed by the European Community Research Infrastructure Action under the FP7 'Capacities' Programme. Paweł Nejfeld and Adam Stebel thank Dr Stanisław Rosadziński (Poznań) for information about the occurrence of *Scutellaria minor* in Lubusz Voivodeship. Researches of Marcin Nobis and Arkadiusz Nowak were financially supported by the National Science Centre, Poland based on the decision no. 2013/09/B/NZ8/03287.

### Disclosure statement

No potential conflict of interest was reported by the authors.

### Notes on contributors

The interests of all the authors lie in taxonomy, chorology and ecology of plants.

**Marcin Nobis:** Contributions the idea and project coordination, field and/or laboratory research, preparing the manuscript

**All others authors:** Contributions field and/or laboratory research and preparing selected parts of the manuscript

### References

- Abdulina, S. A. 1999. *Checklist of Vascular Plants of Kazakhstan*, edited by R. V. Kamelin. Almaty: Ministry of Science, Academy of Sciences of the Republic of Kazakhstan, Institute of Botany and Phytointroduction.
- Alegro, A. L., M. Biljaković, S. Bogdanović, and I. Boršić. 2003. "*Sporobolus pungens* (Schreber) Kunth (Poaceae), rare and endangered psammophytic plant species in Croatia." *Natura Croatica* 12: 1–7.
- Alexeev, Yu E. 1991. "The addition on the flora of Asia Media and Kazakhstan." *Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Biologicheskii* 96 (4): 115–116.
- Amarell, U. 2013. "*Panicum riparium* H. Scholz – eine neoindigene Art Europas? [*Panicum riparium* H. Scholz – a neoindigen species in Europe?]." *Kochia* 6: 1–24.
- Baenitz, G. 1861. *Flora der östlichen Niederlausitz [The flora of eastern Lower Lusatia]*. Görlitz.
- Baikov, K. S., ed. 2005. *Conspectus florae Sibiriae: plantae vasculares*. Novosibirsk: Nauka.
- Baikov, K. S., editor. 2012. *Conspectus florae Rossicae Asiaticae: plantae vasculares*. Novosibirsk: Publishing House of the Siberian Branch of the Russian Academy of Science.
- Bajtenov, M., and I. Kameneckaja. 1990. "Zur Herausbildung von Zwiebeln der Sektion *Oreiprason* aus dem Tjan-Shan [On origin of Alliums of the section *Oreiprason* in Tian Shan]." *Feddes Repertorium* 101: 159–164.
- Balansa, B. 1874. "Catalogue des graminées du Lazistan." *Bulletin de la Société Botanique de France* 21: 10–20.
- Beck von Mannagetta, G. 1930. "Orobanchaceae." In *Das Pflanzenreich Regni Vegetabilis Conspectus*, vol. IV.261, edited by A. Engler, 1–348, Leipzig: Verlag von Wilhelm Engelmann.
- Bobrov, A. A., J. Zalewska-Gałosz, M. Jopek, and E. A. Movergoz. 2015. "*Ranunculus schmalhauseni* (section *Batrachium*, Ranunculaceae), a neglected water crowfoot endemic to Fennoscandia—a case of rapid hybrid speciation in postglacial environment of North Europe." *Phytotaxa* 233: 101–138.
- Bor, N. L. 1960. *Grasses of Burma, Ceylon, India and Pakistan (excluding Bambuseae)*. London: Pergamon Press.
- Bor, N. L. 1968. "Gramineae." In *Flora of Iraq* 9, edited by C. C. Townsend, E. Guest and A. Al-Rawi, 1–588. Baghdad: Ministry of Agriculture of the Republic of Iraq.
- Bor, N. L. 1970. *Flora Iranica* 70: 1–573. Graz: Akademische Druck- u. Verlagsanstalt.
- Bubnova, S.V. 1990. "*Eleocharis* R. Br." In *Flora Sibiri* 3, edited by G.A. Peshkova and L.I. Malyshev, 25–31. Novosibirsk: Nauka.
- Butcher, R. W. 1960. "Notes on water buttercups." *The Naturalist Hull* 1960: 123–125.
- Carlón, L., G. Gómez Casares, M. Láinz, G. Moreno Moral, Ó. Sánchez Pedraja, and G. M. Schneeweiss. 2008. "Más, a propósito de algunas *Phelipanche* Pomel, *Boulardia* F. W. Schultz y *Orobanche* L. (Orobanchaceae) del oeste

- del Paleártico. [The purpose of some *Phelipanche* Pomel, *Boulardia* F. W. Schultz and *Orobanche* L. (Orobanchaceae) from Western Palaearctic]. *Documentos Jardin Botánico Atlántico (Gijón)* 6: 1–128.
- Carlón, L., G. Gomez Casares, M. Lainz, G. Moreno Moral, O. Sánchez Pedraja, and G.M. Schneeweiss. 2015. "Index Orobanchaceae." Available from: <http://www.farmalierganes.com/Otrospdf/publica/Orobanchaceae%20Index.htm> (accessed 22 December 2015).
- Chepinoga, V. V., and S. A. Rosbakh. 2012. "Aquatic vegetation (Lemneta) in Baikal Siberia." *Vegetation of Russia* 21: 106–123.
- Ciardo, F., S. Jutzeler, F. Hoffer-Massard, C. Bornand, editors. 2011. "Notes floristiques vaudoises 2011." *Bulletin du Cercle vaudois de botanique (Lausanne)* 40: 117–147.
- Çiçek, M., and A. E. Yaprak. 2011. "A new natural hybrid of *Scutellaria* (Lamiaceae) from Turkey." *Phytotaxa* 29: 51–55.
- Clayton, W.D. 1980. "*Panicum* L." In *Flora Europaea* 5, edited by T.G. Tutin, V.H. Heywood, N.A. Burges, D.M. Moore et al., 261. Cambridge: Cambridge University Press.
- Clements, D. R., D. DiTommaso, S. J. Darbyshire, P. B. Cavers, and D. Sartono. 2004. "The biology of Canadian weeds. 127. *Panicum capillare* L." *Canadian Journal of Plant Science* 84: 327–341.
- Conert, H. J., ed. 1998. *Gustav Hegi Illustrierte Flora von Mitteleuropa*, 1(3), Spermatophyta: Angiospermae: Monocotyledones 1(2) Poaceae. Berlin: Parey Buchverlag.
- Cook, C. D. K. 1966. "A monographic study of *Ranunculus* subgenus *Batrachium* (DC.) A. Gray." *Mitteilungen der Botanischen Staatssammlung München* 6 (1): 47–237.
- Cope, T. A. 1982. "Poaceae." In *Flora of Pakistan* 143, edited by E. Nasir and S. I. Ali. Karachi: University of Karachi.
- Costa, J. C., C. Neto, C. Aguiar, J. Capelo, M. D. Espírito Santo, J. Honrado, et al. 2012. "Vascular plant communities in Portugal (Continental, Azores and Madeira)." *Global Geobotany* 2: 1–180.
- Decker, P. 1911. "Beiträge zur Flora der Sulichen Neumark und östlichen Niederlausitz [Contribution to the flora of southern Neumark and eastern Lower Lusatia]." *Verhandlungen des Botanischen Vereins der Provinz Brandenburg und die angrenzenden Länder* 53: 87–269.
- Dickoré, W.B. 1995. "Systematische revision und chorologische analyse der Monocotyledoneae des Karakorum (Zentralasien, West-Tibet) [Systematic revision and chorological analysis of Monocotyledoneae of Karakorum (Central Asia, West-Tibet)]. *Flora Karakorumensis* 1. Angiospermae, Monocotyledoneae." *Stapfia* 39: 1–298.
- Doğan, M. 1985. "*Agrostis* L." In *Flora of Turkey and the East Aegean Islands* 9, edited by P. H. Davis, 348–354. Edinburgh: Edinburgh University Press.
- Ebel, A. L. 2012. *Synopsis of the flora of north-west part of Altay-Sayan province*, edited by A. S. Revushkin. Kemerovo: Irbis. [In Russian].
- Egorova, T. V. 1977. "*Allium* L." In *Plantae Asiae Centralis (seculi materis Instituti botanici nomine V.L. Komarovii)* 7, edited by V. I. Grubov, and T. V. Egorova, 18–67. Leningrad: Nauka, Leningradskoye Otdelenie.
- Egorova, T. V. 2001. "Generis *Eleocharis* R. Br. (Cyperaceae) florum Rossiae conspectus systematicus." *Novosti Sistematiki Vysshikh Rastenii* 33: 56–85.
- Egorova, T. V. 2007. "Synopsis taxonomica generis *Eleocharis* R. Br. (Cyperaceae) florum Europae." *Novosti Sistematiki Vysshikh Rastenii* 39: 159–192.
- Ellenberg, H. 1996. *Vegetation Mitteleuropas mit den Alpen in ökologischer, dynamischer und historischer Sicht [Vegetation of Central Europe with the Alps from ecological, dynamic and historical point of view]*. Stuttgart: Ulmer.
- Ellenberg, H., H. E. Weber, R. Düll, V. Wirth, W. Werner, and D. Paulißen. 1992. *Zeigerwerte von Pflanzen in Mitteleuropa [Indicator values for plants in Central Europe]*. 2nd ed. Göttingen: Verlag Erich Goltze KG.
- Fischer, E. 2004. "Scrophulariaceae." In *The Families and Genera of Vascular Plants* 7, edited by J.W. Kadereit: 333–432. Berlin – Heidelberg – New York: Springer Verlag.
- Fitter, A. H., and H. J. Peat. 1994. "The ecological flora database." *Journal of Ecology* 82: 415–425.
- Frajman, B., and P. Schönschetter. 2008. "Notes on some rare *Orobanche* and *Phelipanche* species (Orobanchaceae) in Croatia." *Acta Botanica Croatica* 67 (1): 103–107.
- Freckmann, R. W., and M. G. Lelong. 2007. "*Panicum* L." In *Manual of Grasses for North America*, edited by M. E. Barkworth, L. A. Anderton, K. M. Capels, S. Long and M. B. Piep, 289–296. Logan, Utah: Intermountain Herbarium & Utah State University Press.
- Freitag, H. 1985. "The genus *Stipa* (Gramineae) in southwest and south Asia." *Notes from the Royal Botanic Garden, Edinburgh* 42: 355–489.
- Fürnrohr, F. editor. 2015. "Bemerkenswerte Pflanzenfunde im Regnitzgebiet seit 2002. [Interesting floristic records in Regnitzgebiet since 2002]" *RegnitzFlora – Mitteilungen des Vereins zur Erforschung der Flora des Regnitzgebietes* 7: 72–76.
- Gregor, T. 2003. "*Eleocharis mamillata* – Distribution and infraspecific differentiation." *Folia Geobotanica* 38 (1): 49–64.
- Gubanov, I. A. 1996. *Conspectus of Flora of Outer Mongolia (vascular plants)*. Moscow: Valang.
- Hansen, A. 1980. "*Sporobolus* R. Br." In *Flora Europaea* 5, edited by T.G. Tutin, V.H. Heywood, N.A. Burges, D.M. Moore, et al., 257–258. Cambridge: Cambridge University Press.
- Harms, R. T. 2015. "A survey of the *Sporobolus compositus* and *Sporobolus vaginiflorus* complexes (Poaceae) in Texas." *Phytoneuron* 49: 1–27.
- Hedge, I. 1968. "Brassicaceae." In *Flora Iranica* 57, edited by K. H. Rechinger. Graz: Akademische Druck- u. Verlagsanstalt.
- Hellwig, T. 1897. "Beiträge zur Florenkenntnis der Provinz Posen [Contribution to the knowledge of the flora of the Province of Posen]." *Zeitschrift der Botanischen Abteilung, Naturwissenschaftlicher Verein der Provinz Posen* 4: 1–10.
- Hohla, M. 2006. "*Panicum riparium* (Poaceae) – neu für Österreich – und weitere Beiträge zur Kenntnis der Adventivflora Oberösterreichs. [*Panicum riparium* (Poaceae) – new for Austria – and other records for the non-indigenous flora of Upper Austria]." *Neulreichia* 4: 9–44.
- Hohla, M. 2013. "*Eragrostis amurensis*, *Euphorbia serpens* und *Lepidium latifolium* – neu für Oberösterreich sowie weitere Beiträge zur Flora Österreichs. [*Eragrostis amurensis*, *Euphorbia serpens* und *Lepidium latifolium* – new for Upper Austria – and other additions for the flora of Austria]." *Stapfia* 99: 35–51.
- Hohla, M., W. Diewald, and G. Király. 2015. "*Limonium gmelini* – eine Steppenpflanze an österreichischen Autobahnen sowie weitere Neuigkeiten zur Flora Österreichs. [*Limonium gmelini* – a steppe plant on Austrian highways and other novelties for the flora of Austria]." *Stapfia* 103: 127–150.
- Holmes, N. T. H. 1979. "A guide to the identification of *Batrachium Ranunculus* species of Britain." *Nature Conservancy Council, Chief Scientist's Team Notes* 14: 1–31.
- Hong, Deyuan, H. Yang, J. Cun-li, M. A. Fischer, N. H. Holmgren, and R. R. Mill. 1998. "Scrophulariaceae A.L.



- Jussieu" In *Flora of China* 18, edited by Z. Y. Wu and P. H. Raven, 1–212. Beijing: Science Press and St.-Louis: Missouri Botanical Garden Press.
- Hooker, J. D. 1890. *The Flora of British India*, 5. London: Reeve & Co.
- Horvatić, S., and L. Gospodarić. 1960. "The occurrence of *Sporobolus vaginaeflorus* (Torr.) Wood. in Croatia [in Croatian]." *Acta Botanica Croatica* 18 (19): 79–103.
- Ivanina, L. I. 1991. "Scrophulariaceae Juss." In *Sosudistye rasteniya sovet'skogo Dal'nego Vostoka* [Vascular plants of the Soviet Far East] 5, edited by S. S. Kharkevich, 287–371. St.-Petersburg: Nauka.
- Ivanov, V. V. 1989. *Opređelitel' rasteniy Severnogo Prikaspiya* [Identification manual of the plants of the Northern Caspian Sea]: *Chenopodiaceae* & *Liliaceae*. Leningrad: Nauka. [In Russian].
- Ivleva, V. I. 2010. "Palynomorphological study of Altai representatives of the genus *Geranium* L. (Geraniaceae)." *Turczaninowia* 13 (3): 140–146.
- Jackowiak, B., Z. Celka, J. Chmiel, K. Latowski, and W. Żukowski. 2007. "Red list of vascular flora of Wielkopolska (Poland)." *Biodiversity. Research and Conservation* 5 (8): 95–127.
- Kadereit, G., E. Zacharias, E. Mavrodiev, and A. P. Sukhorukov. 2010. "Molecular phylogeny of *Atripliceae* (Chenopodioideae, Chenopodiaceae): Implications for systematics, biogeography, flower and fruit evolution, and the origin of C4 photosynthesis." *American Journal of Botany* 97 (10): 1664–1687.
- Kapitonova, O. A., V. I. Kapitonov, and N. G. Il'minskikh. 2014. "About the record of the *Utricularia macrorhiza* (Lentibulariaceae) in the West Siberia." *Turczaninowia* 17 (2): 82–86.
- Karelin, G., and J. Kirilow. 1842. "Enumeratio plantarum in desertis Songoriae orientalis et in jugo summorum alpium Alatau anno 1841 collectarum (Continuatio...)." *Bulletin de la Société Impériale des Naturalistes de Moscou* 15: 503–542.
- Khassanov, F. O. 1992. "A revision of the genus *Allium* L. in the flora of Uzbekistan." In *The Genus Allium—Taxonomic Problems and Genetic Resources: Proceedings of an International Symposium held at Gatersleben, Germany, June 11–13, 1991*, edited by P. Hanelt, K. Hammer and H. Knüpffer, 153–159. Gatersleben: Institut für Pflanzengenetik und Kulturpflanzenforschung.
- Király, G., and A. Alegro. 2015. "Re-evaluation of the *Panicum capillare* complex (Poaceae) in Croatia." *Acta Botanica Croatica* 74: 173–179.
- Király, G., A. Baranyai-Nagy, and Sz. Kerekes, A. Király and M. Korda. 2009. "Additions to the knowledge of the alien flora of Hungary [in Hungarian]." *Flora Pannonica* 7: 3–31.
- Király, G., and M. Hohla. 2015. "New stage of the invasion: *Sporobolus vaginiflorus* (Poaceae, Chloridoideae) reached Hungary." *Studia botanica hungarica* 46: 149–155.
- Klimeš, L., and W. B. Dickoré. 2005. "A contribution to the vascular plant flora of Lower Ladakh (Jammu & Kashmir, India)." *Willdenowia* 35: 125–153.
- Kotuhov, J. A., A. N. Danilova, and O. A. Anufrieva. 2011. "Abstract of onions (*Allium* L.) of the Kazakhstan Altai, Sauro-Monrak and Zajsan Hollow." *Botanicheskiye issledovaniya Sibiri i Kazakhstana* 17: 3–33.
- Kurchenko, E. I. 2010. *Genus Agrostis L. (Poaceae) of Russia and neighbouring countries*. Moscow: Promethy.
- Lazkov, G. A., and G. A. Redina. 2007. "On some alien plant species in Kirghizia." *Botanical Journal (St. Petersburg)* 92 (8): 1240–1243. [In Russian].
- Lazkov, G. A., and A. Sennikov. 2014. "New records in vascular plants alien to Kyrgyzstan." *Biodiversity Data Journal* 2: e1018.
- Lazkov, G. A., and B. A. Sultanova. 2011. "Checklist of vascular plants of Kyrgyzstan." *Norrinia* 24: 1–166.
- Lisitsyna, L. I., and V. G. Papchenkov. 2000. *Flora vodoemov Rossii: Opređelitel' sosudistyh rastenij* [Flora of water reservoirs of Russia: Key to determination of vascular plants]. Moscow: Nauka.
- Lomonosova, M. N. 1990. "*Stipa* L." In *Flora Sibiriae, Poaceae (Graminae)* 2, edited by L. I. Malyshev and G. A. Peschkova, 222–230. Divisio Sibirica: Nauka, Novosibirsk.
- Malo, S., and L. Shuka. 2008. "New records on the flora of the Gjirakastra region (South Albania)." *Natura Montenegrina* 7 (3): 369–373.
- Martinovský, J. O. 1980. "*Stipa* L." In *Flora Europaea* 5, edited by T. G. Tutin, V. H. Heywood, N. A. Burges, D. M. Moore, et al., 247–252. Cambridge: Cambridge University Press.
- Medvedeva, N. A. 1996. "*Atriplex*." In *Flora Vostochnoj Evropy* [Flora of the Eastern Europe] 9, edited by N. N. Tzvelev, 44–54. Sankt-Peterburg: Mir & Semya. [In Russian].
- Melzer, H. 2003. "*Sporobolus vaginiflorus* (Poaceae), ein Neubürger aus Nordamerika, lange übersehen in Österreich – und anderes Neue zur Flora von Kärnten. [*Sporobolus vaginiflorus* (Poaceae), a North American non-native species overlooked for a long time in Austria – and other novelties for the flora of Carinthia]." *Neilreichia* 2–3: 131–142.
- Mirek, Z., H. Piękoś-Mirkowa, A. Zając, and M. Zając. 2002. *Flowering plants and pteridophytes of Poland. A checklist*. Kraków: W. Szafer Institute of Botany, Polish Academy of Sciences.
- Moser, H. 1934. "Untersuchungen über die Blattstruktur von *Atriplex*-Arten und ihre Beziehungen zur Systematik" [Investigations of the leaf structure of *Atriplex* species in relation to their systematics]." *Beihefte zum Botanischen Centralblatt* 52B (2): 378–388.
- Nagy, M., G. Király, L. Magyar, L. Nagy, and Z. Simon. 2012. "Distribution and threats of *Panicum riparium* in Hungary." *Agrofórum* 23 (5): 10–18. [in Hungarian].
- Nakhutsrishvili, G. 2013. *The vegetation of Georgia (South Caucasus)*. Berlin/Heidelberg: Springer.
- Nersesian, A. A. 2004. "The genus *Agrostis* L. (Poaceae) in the Armenian flora." *Flora, rastitel'nost' i rastitel'nye resursy Armenii* 15: 19–21.
- Nikitin, V. V., and A. M. Geldykhonov. 1988. *Manual of vascular plants of Turkmenistan*. Leningrad: Science Publishers. [In Russian].
- Nikitina, E. V., and L. I. Kaschenko. 1951. "Rod *Allium* L.—Luk." In *Flora Kirgizskoy SSR* 3, edited by A. I. Vvedensky, 50–96. Frunze: KirgizFAN SSSR.
- Nobis, M. 2011. "Remarks on the taxonomy and nomenclature of the *Stipa tianschanica* complex (Poaceae), on the base of a new record for the flora of Tajikistan (central Asia)." *Nordic Journal of Botany* 29: 194–199.
- Nobis, M. 2012. "*Stipa narynica* sp. nov. (Poaceae) from the western Tian-Shan Mountains." *Nordic Journal of Botany* 30: 70–76.
- Nobis, M. 2013. "Taxonomic revision of the *Stipa lipskyi* group (Poaceae: *Stipa* section *Smirnovia*) in the Pamir Alai and Tian-Shan Mountains." *Plant Systematics and Evolution* 299: 1307–1354.
- Nobis, M. 2014. "Taxonomic revision of the Central Asian *Stipa tianschanica* complex (Poaceae) with particular reference to the epidermal micromorphology of the lemma." *Folia Geobotanica* 49: 283–308.
- Nobis, M., A. L. Ebel, A. Nowak, B. Paszko, A. A. Bobrov, Y. A. Kotukhov, A. N. Kupriyanov, et al. 2015. "Contribution to the flora of Asian and European countries: new national and regional vascular plant records, 4." *Acta Botanica Gallica: Botany Letters* 162 (4): 301–316.

- Nobis, M., A. L. Ebel, A. Nowak, O. T. Turginov, A. N. Kupriyanov, A. Nobis, et al. 2014. "Contribution to the flora of Asian and European countries: new national and regional vascular plant records, 2." *Acta Botanica Gallica: Botany Letters* 161 (2): 209–221.
- Nobis, M., and P. D. Gudkova. 2016. "Taxonomic notes on feather grasses (Poaceae: *Stipa*) from eastern Kazakhstan with typification of seven names and one new combination." *Phytotaxa* 245 (1): 31–42.
- Nobis, M., E. Klichowska, A. Nowak, P. D. Gudkova, and K. Rola. 2016. "Multivariate morphometric analysis of the *Stipa turkestanica* group (Poaceae)." *Plant Systematics and Evolution* 302 (2): 137–153.
- Nobis, M., A. Nobis, A. Nowak, and S. Nowak. 2014. "*Stipa klimesii* (Poaceae) a new species from Western Himalayas (India)." *Phytotaxa* 174 (3): 173–180.
- Nobis, M., A. Nowak, A. Nobis, B. Paszko, R. Piowarczyk, S. Nowak, et al. 2014. "Contribution to the flora of Asian and European countries: new national and regional vascular plant records." *Acta Botanica Gallica: Botany Letters* 161: 81–89.
- Nobis, M., A. Nowak, A. L. Ebel, A. Nobis, S. Nowak, P. D. Gudkova, et al. 2015. "Contribution to the flora of Asian and European countries: new national and regional vascular plant records, 3." *Acta Botanica Gallica: Botany Letters* 162 (2): 103–115.
- Novopokrovskij, I. V., and N. N. Tzvelev. 1958. "Orobanchaceae." In *Flora Unionis Republicarum Socialistarum Sovieticarum (Flora URSS)*, vol. 23, 685–687, edited by B. K. Shishkin, Institutum Botanicum nomine V. L. Komarovii Academiae Scientiarum URSS, Mosqua – Leningrad.
- Ochyra, R., J. Żarnowiec, and H. Bednarek-Ochyra. 2003. *Census catalogue of Polish mosses*. Kraków: W. Szafer Institute of Botany, Polish Academy of Sciences.
- Olonova, M. V. 1996. "Lentibulariaceae." In *Flora Sibiri* 12, edited by A. V. Polozhij and G. A. Peschkova, 99–102. Novosibirsk: Nauka (Siberian Branch).
- Paszko, B. 2014a. "*Agrostis schischkini*, a new name for *Agrostis trichantha* (Schischk.) Tzvelev (Poaceae, Agrostidinae)." *Phytotaxa* 170: 136–138.
- Paszko, B. 2014b. "*Agrostis pendryi* (Poaceae: Agrostidinae)—a new species from the Central Himalaya." *Phytotaxa* 175: 29–36.
- Paszko, B., and C. A. Pendry. 2013. "*Agrostis griffithiana* (Poaceae: Agrostidinae)—typification, a new synonym and an update of the distribution in India." *Phytotaxa* 140: 26–34.
- Paul, T. K. 2012. "A synopsis of the family Chenopodiaceae in India." *Pleione* 6 (2): 273–297.
- Pavlov, N. V., and P. P. Poljakov. 1958. "Rod Luk – *Allium* L." In *Flora Kazakhstana* 2, edited by N. V. Pavlov, 134–193. Alma-Ata: Izdatelstvo AN KazSSR.
- Pazdnikova, N. M., and V. V. Chepinoga. 2013. "Check-list of the vascular flora of the Onon-Dahuria region, central part (South-East Transbaikalia, Russia)." *The Bulletin of Irkutsk State University. Series «Biology, Ecology»* 6(1): 32–60.
- Pazij, V. K. 1968. "*Stipa*." In: *Opredelitel rastenii srednei Azii [Conspectus florae Asiae Mediae] 1*, edited by S. S. Kovalevskaya, 64–82 + 200–201. Taschkent: USSR Academy of Sciences.
- Peschkova, G. A. 1996. "Geraniaceae." In *Flora Sibiri* 10, edited by G. A. Peschkova, 8–23. Novosibirsk: Nauka (Siberian Branch).
- Peterson, P. M., S. L. Hatch, and A. S. Weakley. 2007. "*Sporobolus* R. Br." In *Manual of Grasses for North America*, edited by M. E. Barkworth, L. A. Anderton, K. M. Capels, S. Long, and M. B. Piep, 212–217. Logan: Intermountain Herbarium and Utah State University Press.
- Piowarczyk, R. 2012. "Revised distribution and phytosociological data of *Orobanche coerulescens* Stephan in Willd. (Orobanchaceae): Poland in relation to Central Europe." *Biodiversity Research and Conservation* 26: 61–72.
- Piowarczyk, R. 2015a. "*Orobanche zajaciorum* (Orobanchaceae): a new species from the Caucasus." *Phytotaxa* 201 (3): 214–220.
- Piowarczyk, R. 2015b. "Seed micromorphology of central European *Orobanche* and *Phelipanche* (Orobanchaceae) in relation to preferred hosts and systematic implications." *Australian Systematic Botany* 28: 124–136.
- Piowarczyk, R., and A. Przemyski. 2009. "New locality of *Orobanche coerulescens* Stephan ex Willd. (Orobanchaceae) at the NW limit of its geographical range." *Acta Societatis Botanicorum Poloniae* 78: 291–295.
- Piowarczyk, R., D. Kwolek, M. Denysenko, M. Cygan, G. Góralski, H. Ślesak, et al. 2015. "*Orobanche grenieri* (Orobanchaceae), a southwestern European species newly found in Asia." *Annales Botanici Fennici* 52: 411–418.
- Piowarczyk, R., J. Madeja, and M. Nobis. 2015. "Pollen morphology of the Central European broomrapes (Orobanchaceae: *Orobanche*, *Phelipanche* and *Orobanchella*) and its taxonomical implications." *Plant Systematics and Evolution* 301: 795–808.
- Poljakov, P. P. 1958. "Cyperaceae J. St.-Hil." In *Flora Kazakhstana* 2, edited by N. V. Pavlov, 3–83. Almaty: Akademia nauk Kazakhskoi SSR.
- Pusch, J. 2009. "Orobanchaceae (Sommerwurzgewächse)." In *Illustrierte Flora von Mitteleuropa (Illustrated flora of the central Europe)*, Bd. 6/1A, Lieferung 1, edited by G. Hegi, 1–99. Jena: Weissdorn-Verlag.
- Rätzel, S., and H. Uhlich. 2004. "*Orobanche benkertii* sp. nov. (Orobanchaceae Vent.) und weitere *Orobanche*-Sippen aus dem NW-Kaukasus. [*Orobanche benkertii* sp. nov. (Orobanchaceae Vent.) and other *Orobanche* species from the northwestern Caucasus]." *Feddes Repertorium* 115: 189–211.
- Rozhevitz, R. Y. 1934. "*Poa* L." In *Flora SSSR* 2, edited by V. L. Komarov, 366–426. Leningrad: Izdatelstvo Akademii Nauk SSSR.
- Ryves, T. B., E. J. Clement, and M. C. Foster. 1996. *Alien grasses of the British Isles*. London: BSBI.
- Rzazade, R. Y. 1950. "*Agrostis* L." In *Flora Azerbaydzhan* 1, edited by I. I. Karyagin, 182–186. Baku: Izdatel'stvo Akademii Nauk Azerbajdzanskoj SSR. (in Russian).
- Scholz, H. 2002. "*Panicum riparium* H. Scholz – eine neue indigene Art der Flora Mitteleuropas. [*Panicum riparium* H. Scholz – a new native species to the Central European flora]." *Feddes Repertorium* 113: 273–280.
- Seregin, A. P., G. Anačkov, and N. Friesen. 2015. "Molecular and morphological revision of the *Allium saxatile* group (Amaryllidaceae): geographical isolation as the driving force of underestimated speciation." *Botanical Journal of the Linnean Society* 177 (1): 67–101.
- Seregin, A. P., and N. Friesen. 2015. "Molecular and morphological revision reveals different evolution patterns in *Allium* sect. *Oreiprason* and sect. *Falcatifolia* (Amaryllidaceae)." In *Semicentenary after Konstantin Meyer: XIII Moscow symposium on plant phylogeny: Proceedings of the international conference (February 2–6, 2015, Moscow)*, edited by A. C. Timonin, 353–356. Moscow: MAKSS Press.

- Smith, S. G., J. J. Bruhl, M. S. Gonzales-Elizondo, and F. J. Menapace. 2002. "*Eleocharis* R. Brown." In *Flora of North America* 23, edited by Flora of North America Editorial Committee, 60–121. New York – Oxford: Oxford University Press.
- Starchenko, V. M. 2008. *Flora of Amur Region and Problems of its Conservation: Far East of Russia*. Moscow: Nauka.
- Sukhorukov, A. P. 2005. *Konspekt vidov sosudistykh rastenii Djanybeksogo biologicheskogo stacionara I ego okrestnostej [Checklist of the vascular plants of Djanybek botanical reserve and its surroundings]*. Moscow: Maks-Press. [In Russian].
- Sukhorukov, A. P. 2006. "Zur Systematik und Chorologie der in Russland und benachbarten Staaten (in den Grenzen der ehemaligen UdSSR) vorkommenden *Atriplex*-Arten (*Chenopodiaceae*)." [Systematics and Chorology of the *Atriplex* Species in Russia and Neighboring States] *Annalen des Naturhistorischen Museums in Wien* 108B: 307–420. [In German].
- Sukhorukov, A. P. 2014. *The carpology of the Chenopodiaceae with reference to the phylogeny, systematics and diagnostics of its representatives*. Tula: Grif & Co.. [In Russian with English summary].
- Sukhorukov, A. P., and M. A. Kushunina. 2015. "Corrigenda to "Taxonomic revision of *Chenopodiaceae* in Nepal" [Phytotaxa 191: 10–44. 2014]." *Phytotaxa* 226 (3): 288–291.
- Swan, G. A. 2007. "The *Eleocharis mamillata* H. Lindb. fil. aggregate (Cyperaceae) in the British Isles." *Watsonia* 26: 317–325.
- Szulczewski, J. W. 1951. "Wykaz roślin naczyniowych w Wielkopolsce dotąd stwierdzonych [List of species since this time in the Greater Poland ascertained]." *Prace Komisji Biologicznej Poznańskiego Towarzystwa Przyjaciół Nauk* 12 (6): 1–128.
- Takhtajan, A. L., ed. 1987. *Flora Armenii (Flora of Armenia), Verbenaceae - Lentibulariaceae*, 8. Erevan: Izdatel'stvo Akademii Nauk Armjanskoj SSR.
- Taxonomic *Allium* Reference Collection. 2015. "Facilitated by IPK Gatersleben". Accessed 6 October 2015. <http://apex.ipk-gatersleben.de/apex/f?p=265:1>
- Taylor, P. 1989. "The genus *Utricularia* – a taxonomical monograph." *Kew Bulletin Additional Series* 14: 1–724.
- Teryokhin, E. S., G. B. Schibakina, N. B. Serafimovitsch, and T. I. Kravtsova. 1993. *Opredelitel Sarasychovykh Flory SSSR [Determinator of broomrapes of the USSR flora]*. Sankt Petersburg: Nauka.
- The Plant List. 2013. Version 1.1.: online database. (Accessed 4 March 2016) <http://www.theplantlist.org/>
- Tinner, U. 2013. "Zwei neue Grasarten im St. Galler Rheintal: *Sporobolus vaginiflorus* und *Sporobolus neglectus*." *Bauhinia* 24: 53–56.
- Tokarska-Guzik, B., Z. Dajdok, M. Zając, A. Zając, A. Urbisz, W. Danielewicz, et al. 2012. *Rośliny obcego pochodzenia w Polsce ze szczególnym uwzględnieniem gatunków inwazyjnych [Alien plants in Poland with particular reference to invasive species]*. Warszawa: Generalna Dyrekcja Ochrony Środowiska.
- Troshkina, V. I. 2015. "Type specimens of names of taxa of *Geranium* L. (Geraniaceae) described by L. P. Sergievskaja and deposited in the Krylov Herbarium (TK)." *Novosti Sistematiki Vysshikh Rastenii* 46: 119–125.
- Tzvelev, N. N. 1974. "Zametki o tribe Stipeae Dum. semejstva zlakov (Poaceae) v SSSR –Notulae de tribu Stipeae Dum. (fam. Poaceae) in URSS." *Novosti Sistematiki Vysshikh Rastenii* 11: 4–21.
- Tzvelev, N. N. 1976. *Zlaki SSSR [Grasses of the Soviet Union]*. Leningrad: Nauka.
- Tzvelev, N. N. 2006. "Fam. 178. *Poaceae* Barnhart (Gramineae Juss.)." In *Caucasian Flora Conspectus* 2, edited by A.L. Takhtajan, 248–378. St Petersburg: St. Petersburg University Press (in Russian).
- Tzvelev, N. N. 2012. "Notes on the tribe Stipeae Dumort. (Poaceae)." *Novosti Sistematiki Vysshikh Rastenii* 43: 20–29.
- Urbisz, A. 2011. "Occurrence of temporarily-introduced alien plant species (ephemerophytes) in Poland - scale and assessment of the phenomenon." *Prace Naukowe Uniwersytetu Śląskiego w Katowicach* 2897: 1–198.
- Urgamal, M., B. Oyuntsetseg, D. Nyambayar, and Ch. Dulamsuren. 2014. *Conspectus of the Vascular Plants of Mongolia*. Ulaanbaatar: 'Admon' Press.
- Verloove, F. 2014. "Manual of the Alien Plants of Belgium. *Panicum capillare*." Accessed 21 January 2014. <http://alienplantsbelgium.be>
- Vvedensky, A. I. 1941. "Rod *Allium* L.—Luk. [Genus *Allium* L.—Leek.]" In *Flora Uzbekistana* 1, edited by R. R. Shreder, 427–467. Tashkent: UzFAN SSSR.
- Vvedensky, A. I. 1971. "Rod *Allium* L.—Luk. [Genus *Allium* L.—Leek.]" In *Opredelitel rasteniy Sredney Azii: Kriticheskiy konspekt flory* 2, edited by S. S. Kovalevskaya, 39–89 & 311–328. Tashkent: FAN Uzbekskoy SSR.
- Walters, S. M. 1980. "*Eleocharis* R. Br." In *Flora Europaea* 5, edited by T. G. Tutin, V. H. Heywood, N. A. Burges, and D. H. Valentine, 281–284. Cambridge University Press, Cambridge.
- Webster, S. D. 1988. "*Ranunculus penicillatus* (Dumort.) Bab. in Great Britain and Ireland." *Watsonia* 17: 1–22.
- Wilhelm, T. 1998. "Neues zur Ausbreitung von *Sporobolus vaginiflorus* (Torrey) Wood." *Floristische Rundbriefe* 32: 95–100.
- Wilhelm, T. 2011. "Ergänzungen und Korrekturen zum Katalog der Gefäßpflanzen (4)." *Gredleriana* 11: 71–82.
- Wu, Z. L., and S. M. Phillips. 2006. "Tribe Stipeae." In *Flora of China: Poaceae* 22, edited by Z. Y. Wu et al., 188–212. Beijing: Science Press & St Louis: Missouri Botanical Garden Press.
- Xu, J. M., and R. V. Kamelin. 2000. "*Allium* Linnaeus." In *Flora of China* 24, edited by Z. Y. Wu and P. H. Raven, 165–202. Beijing: Science Press; St Louis: Missouri Botanical Garden Press.
- Yunusov, S. Yu. 1978. "Katran – *Crambe* L." In *Flora of Tajikistan* 5, edited by P. N. Ovchinnikov, 222–224. Leningrad: Nauka.
- Zalewska-Gałosz, J., M. Jopek, and T. Ilnicki. 2015. "Hybridization in *Batrachium* group: Controversial delimitation between heterophyllous *Ranunculus penicillatus* and the hybrid *Ranunculus fluitans* × *R. peltatus*." *Aquatic Botany* 120: 160–168.
- Zazanashvili, N., R. Gagnidze, and G. Nakhutsrishvili. 2000. "Main types of vegetation zonation on the mountains of the Caucasus." *Acta Phytogeographica Suecica* 85: 7–16.
- Zhu (Chu), G.-L., S. L. Mosyakin, and S. E. Clemants. 2003. "*Chenopodiaceae*." In *Flora of China* 5. *Ulmaceae–Basellaceae*, edited by Z. Wu, and P. H. Raven, 351–414. Beijing: Science Press.