

Free-Radical Nonbranched-Chain Oxidation of Hydrogen

Michael M. Silaev

Abstract—New reaction scheme is suggested for the initiated nonbranched-chain addition of hydrogen atoms to the multiple bond of the molecular oxygen. The scheme includes the addition reaction of the hydroperoxyl free radical to the oxygen molecule to form the hydrotetraoxyl free radical which is relatively low-reactive and inhibits the chain process by shortening of the kinetic chain length. This reaction competes with chain propagation reactions through a hydrogen atom. Based on the proposed scheme rate equations (containing one to three parameters to be determined directly) are deduced using quasi-steady-state treatment. The kinetic description with use the obtained rate equations is applied to the γ -induced nonbranched-chain process of the free-radical oxidation of hydrogen dissolved in water containing different amounts of oxygen at 296 K. The ratio of rate constants of competing reactions and the rate constant of the addition reaction to the molecular oxygen are defined. In this process the oxygen with the increase of its concentration begins to act as an oxidation autoinhibitor (or an antioxidant), and the rate of hydrogen peroxide formation as a function of the dissolved oxygen concentration has a maximum. From the energetic standpoint possible nonchain pathways of the free-radical oxidation of hydrogen and the routes of ozone decay *via* the reaction with the hydroxyl free radical (including the addition yielding the hydrotetraoxyl free radical) in the Earth's upper atmosphere were considered.

Index Terms—competition, hydrogen, low-reactive hydrotetraoxyl free radical, thermochemical data, energy.

I. INTRODUCTION

The kinetics of inhibition for nonbranched-chain processes of saturated free-radical addition to the C=C and C=O double bonds of alkene and formaldehyde molecules, respectively, by low-reactive free radicals that can experience delocalization of the unpaired p -electron was first considered in [1]. In these processes a low-reactive free radical is formed in the reaction competing

with chain propagation reactions through a reactive free radical. In the present work the kinetics of inhibition by low-reactive hydrotetraoxyl free radical is considered for the nonbranched-chain process of the addition of a hydrogen atom to one of the two multiply bonded atoms of the oxygen molecule yielding a hydroperoxyl free radical. The hydroperoxyl free radical then abstracts the most labile atom from a molecule of the compound being oxidized or decomposes to turn into a molecule of an oxidation product. The only reaction that can compete with these two reactions at the chain evolution stage is the addition of the peroxy radical to the oxygen molecule (provided that the oxygen concentration is sufficiently high). This reaction yields the secondary hydrotetraoxyl 1:2 adduct radical, which is the heaviest and the largest among the reactants. It is less reactive than the primary peroxy 1:1 adduct radical and, as a consequence, does not participate in further chain propagation. At moderate temperatures, the reaction proceeds *via* a nonbranched-chain mechanism.

The aim of this study was the mathematical simulation of oxidation process autoinhibited by oxygen, when the dependence of the peroxide formation rate on the dissolved oxygen concentration has a maximum. The simulation was based on experimental data obtained for γ -radiation-induced addition reaction of hydrogen atom to the molecular oxygen for which the initiation rate V_1 is known (taking into account that $V = GP$ and $V_1 = G(H^*)P$, where P is the dose rate, and $G(H^*)$ is the initial yield of the chain-carrier hydrogen atom H^* – initiation yield [2, 3]).

Based on the reaction scheme suggested for the kinetic description of the addition process to oxygen, the kinetic equations with one to three parameters to be determined directly were derived. Reducing the number of unknown parameters in a kinetic equation will allow one to decrease the narrowness of the correlation of these parameters and to avoid a sharp buildup of the statistical error in the nonlinear estimation of these parameters in the case of a limited number of experimental data points. The rate constant of the addition to oxygen, estimated as a kinetic parameter, can be compared to its reference value if the latter is known. This provides a clear criterion to validate the mathematical description against experimental data.

II. KINETICS OF HYDROGEN ATOM ADDITION TO THE OXYGEN MOLECULE

A number of experimental findings concerning the autoinhibiting effect of an increasing oxygen concentration at modest temperatures on hydrogen

Manuscript Received August 20, 2018

Michael M. Silaev, Chemistry Faculty, Lomonosov Moscow State University, Vorobievsky Gory, Moscow, 119991, Russia, (e-mail: mmsilaev@rc.chem.msu.ru).

oxidation both in the gas phase [4–6] (Fig. 2) and in the liquid phase [7] (Fig. 1, curve 2), considered in my earlier works [8–11], can also be explained in terms of the competition kinetics of free radical addition [12, 13]. From Fig. 2 shows that the quantum yields of hydrogen peroxide and water (of products of photochemical oxidation of hydrogen at atmospheric pressure and room temperature) are maximum in the region of small concentrations of oxygen in the hydrogen–oxygen system (curves 1 and 2, respectively) [4].

In the familiar monograph “Chain Reactions” by Semenov [14], it is noted that raising the oxygen concentration when it is already sufficient usually slows down the oxidation process by shortening the chains. The existence of the upper (second) ignition limit in oxidation is due to chain termination in the bulk through triple collisions between an active species of the chain reaction and two oxygen molecules (at sufficiently high oxygen partial pressures). In the gas phase at atmospheric pressure, the number of triple collisions is roughly estimated to be 10^3 times smaller than the number of binary collisions (and the probability of a reaction taking place depends on the specificity of the action of the third particle) [14]. Note that in the case of a gas-phase oxidation of hydrogen at low pressures of 25–77 Pa and a temperature of 77 K [5] when triple collisions are unlikely, the dependence of the rate of hydrogen peroxide formation on oxygen concentration (the rate of passing of molecular oxygen *via* the reaction tube) also has a pronounced maximum (see curves 3 and 4 in Fig. 2) that indicates a chemical mechanism providing the appearance of a maximum (see reaction 4 of *Scheme*).




Figure 1. (1, □) Reconstruction of the functional dependence of the total hydrogen peroxide formation rate $V_3, \gamma(\text{H}_2\text{O}_2)$ on the dissolved oxygen concentration x from empirical data (symbols) using Eqs. (1a) and (4a) (model optimization with respect to the parameter α) for the γ -radiolysis of water saturated with hydrogen and containing different amounts of oxygen at 296 K [7] ($S_Y = 1.13 \times 10^{-8}$). (2) The dashed curve described $V_3(\text{H}_2\text{O}_2)$ as a function of the oxygen concentration x based on Eq. (1a) (model optimization with respect to α) and the experimental data of curve 2 ($S_Y = 1.73 \times 10^{-8}$).




Figure 2. (1, 2) Quantum yields of (1, ●) hydrogen peroxide and (2, ○) water resulting from the photochemical oxidation of hydrogen in the hydrogen–oxygen system as a function of the oxygen concentration x (light wavelength of 171.9–172.5 nm, total pressure of 10^5 Pa, room temperature [4]). (3, 4) Hydrogen peroxide formation rate $V(\text{H}_2\text{O}_2)$ (dashed curves) as a function of the rate $V(\text{O}_2)$ at which molecular oxygen is passed through a gas-discharge tube filled with (3, △) atomic and (4, □) molecular hydrogen. Atomic hydrogen was obtained from molecular hydrogen in the gas-discharge tube before the measurements (total pressure of 25–77 Pa, temperature of 77 K [5]). The symbols represent experimental data.

Scheme



Nonbranched-chain oxidation of hydrogen and changes in enthalpy (ΔH_{298}° , kJ mol $^{-1}$) for elementary reactions¹

Chain initiation




¹According to Francisco and Williams [15], the enthalpy of formation (ΔH_f°) in the gas phase of H^\cdot , HO^\cdot , HO_2^\cdot , HO_4^\cdot (the latter without the possible intramolecular hydrogen bond taken into account), O_3 , H_2O [6], H_2O_2 , and H_2O_4 is 218.0 ± 0.0 , 39.0 ± 1.2 , 12.6 ± 1.7 , 122.6 ± 13.7 , 143.1 ± 1.7 , -241.8 ± 0.0 , -136.0 ± 0 , and -26.0 ± 9 kJ mol $^{-1}$, respectively. Calculations for the HO_4^\cdot radical with a helical structure were carried out using the G2(MP2) method [16]. The stabilization energies of HO_2^\cdot , HO_4^\cdot , and HO_3^\cdot were calculated in the same work to be 64.5 ± 0.1 , 69.5 ± 0.8 , and 88.5 ± 0.8 kJ mol $^{-1}$, respectively. The types of the O_4 molecular dimers, their IR spectra, and higher oxygen oligomers were reported [17, 18]. The structure and IR spectrum of the hypothetical cyclotetraoxygen molecule O_4 , a species with a high energy density, were calculated by the CCSD method, and its enthalpy of formation was estimated [19]. The photochemical properties of O_4 and the van der Waals nature of the O_2 – O_2 bond were investigated [20, 21]. The most stable geometry of the dimer is two O_2 molecules parallel to one another. The O_4 molecule was identified by NR mass spectrometry [22].




Chain propagation



Inhibition



Chain termination



The chain evolution (propagation and inhibition) stage of the *Scheme* includes consecutive reaction pairs 2–3 and 3–3'; the parallel (competing) reaction pair 3–4; and consecutive–parallel reactions 2 and 4.

The hydroperoxyl free radical HO_2^\bullet [23–26] resulting from reaction 2 possesses an increased energy due to the energy released the conversion of the O=O multiple bond into the HO–O[•] ordinary bond. Therefore, before its possible decomposition, it can interact with a hydrogen or oxygen molecule as the third body *via* parallel (competing) reactions 3 and 4, respectively. The hydroxyl radical HO^\bullet that appears and disappears in consecutive parallel reactions 3 (first variant) and 3' possesses additional energy owing to the exothermicity of the first variant of reaction 3, whose heat is distributed between the two products. As a consequence, this radical has a sufficiently high reactivity not to accumulate in the system during these reactions, whose rates are equal ($V_3 = V_{3'}$) under quasi-steady-state conditions, according to the above scheme. Parallel reactions 3 (second, parenthesized variant) and 3' regenerate hydrogen atoms. It is assumed [9, 10] that the hydrotetraoxyl radical HO_4^\bullet (first reported in [27–29]) resulting from endothermic reaction 4, which is responsible for the peak in the experimental rate curve (Fig. 2, curve 3), is closed into a five-membered $[\text{OO}-\text{H}\cdots\text{OO}]^\bullet$ cycle due to weak intramolecular hydrogen bonding [30, 31]. This structure imparts additional stability to this radical and makes it least reactive.

The HO_4^\bullet radical was discovered by Staehelin *et al.* [32] in a pulsed radiolysis study of ozone degradation in water; its UV spectrum with an absorption maximum at 260 nm ($\varepsilon(\text{HO}_4^\bullet)_{280\text{nm}} = 320 \pm 15 \text{ m}^2 \text{ mol}^{-1}$) was reported.

The spectrum of the HO_4^\bullet radical is similar to that of

ozone, but the molar absorption coefficient $\varepsilon(\text{HO}_4^\bullet)_{\lambda_{\text{max}}}$ of the former is almost two times larger [32]. The assumption about the cyclic structure of the HO_4^\bullet radical can stem from the fact that its mean lifetime in water at 294 K, which is $(3.6 \pm 0.4) \times 10^{-5} \text{ s}$ (as estimated [11] from the value of $1/k$ for the monomolecular decay reaction $\text{HO}_4^\bullet \xrightarrow{k} \text{HO}_2^\bullet + \text{O}_2$ [32]), is 3.9 times longer than that of the linear HO_3^\bullet radical [16, 33] estimated in the same way [11] for the same conditions [34], $(9.1 \pm 0.9) \times 10^{-6} \text{ s}$.

MP2/6-311++G** calculations using the Gaussian-98 program confirmed that the cyclic structure of HO_4^\bullet [35] is energetically more favorable than the helical structure [16] (the difference in energy is 4.8–7.3 kJ mol⁻¹, depending on the computational method and the basis set).² For example, with the MP2(full)/6-31G(d) method, the difference between the full energies of the cyclic and acyclic HO_4^\bullet conformers with their zero-point energies (ZPE) values taken into account (which reduces the energy difference by 1.1 kJ mol⁻¹) is –5.1 kJ mol⁻¹ and the entropy of the acyclic-to-cyclic HO_4^\bullet transition is

$\Delta S_{298}^\circ = -1.6 \text{ kJ mol}^{-1} \text{ K}^{-1}$. Therefore, under standard conditions, HO_4^\bullet can exist in both forms, but the cyclic structure is obviously dominant (87%, $K_{eq} = 6.5$) [35].

Reaction 4 and, to a much lesser degree, reaction 6 inhibit the chain process, because they lead to inefficient consumption of its main participants – HO_2^\bullet and H^\bullet .

The hydrogen molecule that results from reaction 5 in the gas bulk possesses an excess energy, and, to acquire stability within the approximation used in this work, it should have time for deactivation *via* collision with a particle M capable of accepting the excess energy [37]. To simplify the form of the kinetic equations, it was assumed that the rate of the bimolecular deactivation of the molecule substantially exceeds the rate of its monomolecular decomposition, which is the reverse of reaction 5 [38].

Reactions³ 6 and 7 regenerate hydrogen and oxygen (in the form of $\text{O}_2(X^3\Sigma_g^-)$ molecules, including the singlet states with $\Delta H_{f298}^\circ(\text{O}_2, a^1\Delta_g) = 94.3 \text{ kJ mol}^{-1}$ [15, 18]

²There were calculations for the two conformers (*cis* and *trans*) of the HO_4^\bullet radical [36] using large scale *ab initio* methods and density functional techniques with extended basis sets. Both conformers have a nearly planar geometry with respect to the four oxygen atoms and present an unusually long central O–O bond. The most stable conformer of HO_4^\bullet radical is the *cis* one, which is computed to be endothermic with respect to $\text{HO}_2^\bullet(X^2A'') + \text{O}_2(X^3\Sigma_g^-)$ at 0 K.

³Taking into account the principle of detailed balance for the various pathways of formation of products, whose numbers in the elementary reaction should not exceed three for possible involvement in the triple collisions in the case of the reverse reaction, since the probability of simultaneous interaction of four particles is negligible.

and $\Delta H_{f298}^{\circ}(\text{O}_2, b^1\Sigma_g^+) = 161.4 \text{ kJ mol}^{-1}$ [18], which are deactivated by collisions, and in the form of O_3) and yield hydrogen peroxide or water *via* a nonchain mechanism, presumably through the intermediate formation of the unstable hydrogen tetraoxide molecule H_2O_4 [39, 40].⁴ Ozone does not interact with molecular hydrogen. At moderate temperatures, it decomposes fairly slowly, particularly in the presence of $\text{O}_2(X^3\Sigma_g^-)$ [18]. The reaction of ozone with H^{\bullet} atoms, which is not impossible, results in their replacement with HO^{\bullet} radicals. The relative contributions from reactions 6 and 7 to the process kinetics can be roughly estimated from the corresponding enthalpy increments (*Scheme*).

When there is no excess hydrogen in the hydrogen–oxygen system and the homomolecular dimer O_4 [19–22, 41, 42], which exists at low concentrations (depending on the pressure and temperature) in equilibrium with O_2 [18], can directly capture the H^{\bullet} atom to yield the heteronuclear cluster HO_4^{\bullet} ,⁵ which is more stable than O_4 [18] and cannot abstract a hydrogen atom from the hydrogen molecule, nonchain hydrogen oxidation will occur to give molecular oxidation products *via* the disproportionation of free radicals.

The low-reactive hydrotetraoxyl radical HO_4^{\bullet} [32], which presumably has a high energy density [19], may be an intermediate in the efficient absorption and conversion of biologically hazardous UV radiation energy the Earth upper atmosphere. The potential energy surface for the atmospheric reaction $\text{HO}^{\bullet} + \text{O}_3$, in which the adduct $\text{HO}_4^{\bullet}(^2A)$ was considered as an intermediate, was calculated by the DMBE method [43]. From this standpoint, the following reactions are possible in the upper troposphere, as well as in the lower and middle stratosphere, where most of the ozone layer is situated (altitude of 16–30 km, temperature of 217–227 K, pressure of 1.0×10^4 – 1.2×10^3 Pa [44]; the corresponding ΔH_{298}° reaction values are given in kJ mol^{-1} [15]):

8. $\text{H}_2\text{O}(\text{vapor}) + h\nu \rightarrow \text{H}^{\bullet} + \text{HO}^{\bullet}$ [44]
9. $\text{HO}^{\bullet} + \text{O}_3 \rightarrow \text{HO}_4^{\bullet}$ [29, 32, 43], $\Delta H_{298}^{\circ} = -59.5$;
10. $\text{HO}_4^{\bullet} \rightarrow \text{HO}_2^{\bullet} + \text{O}_2(X^3\Sigma_g^-)$ [32, 43], $\Delta H_{298}^{\circ} = -110.0$
(or $\text{HO}_2^{\bullet} + \text{O}_2(a^1\Delta_g)$), $\Delta H_{298}^{\circ} = -15.7$.

The HO_4^{\bullet} radical can disappear *via* disproportionation with a molecule, free radical, or atom in addition to

dissociation. Note that emission from $\text{O}_2(a^1\Delta_g)$ and $\text{O}_2(b^1\Sigma_g^+)$ is observed at altitudes of 30–80 and 40–130 km, respectively [45].

Staehelin *et al.* [32] pointed out that, in natural systems in which the concentrations of intermediates are often very low, kinetic chains in chain reactions can be very long in the absence of scavengers since the rates of the chain termination reactions decrease with decreasing concentrations of the intermediates according to a quadratic law, whereas the rates of the chain propagation reactions decrease according to a linear law.

The kinetic description of the noncatalytic oxidation of hydrogen, including in an inert medium [37], in terms of the simplified scheme of free-radical nonbranched-chain reactions (*Scheme*), which considers only quadratic-law chain termination and ignores the surface effects [5], at moderate temperatures and pressures, in the absence of transitions to unsteady-state critical regimes, and at a substantial excess of the hydrogen concentration over the oxygen concentration was obtained by means of quasi-steady-state treatment, as in the previous studies on the kinetics of the branched-chain free-radical oxidation of hydrogen [46], even though the applicability of this method in the latter case under unsteady states conditions was insufficiently substantiated. The method was used with the following condition⁶ for the first stages of the process: $k_6 = \sqrt{2k_5 2k_7}$ [47] and, hence, $V_1 = V_5 + 2V_6 + V_7 = (\sqrt{2k_5}[\text{H}^{\bullet}] + \sqrt{2k_7}[\text{HO}_4^{\bullet}])^2$ which allow the exponent of the $2k_5[\text{H}^{\bullet}]^2$ term in the $d[\text{H}^{\bullet}]/dt = 0$ equation to be reduced to 1 [1, 47]. The kinetic equations were derived for the rates ($\text{mol dm}^{-3} \text{ s}^{-1}$) of the elementary reactions for the formation of molecular products of hydrogen oxidation.

The kinetic equations were derived for the rates ($\text{mol dm}^{-3} \text{ s}^{-1}$) of the elementary reactions for the formation of molecular products of hydrogen oxidation.

The rate of the chain formation of hydrogen peroxide in propagation reaction 3 and of water in reactions 3 and 3' with $V_{3,3}(\text{H}_2\text{O}) = 2V_3$ is

$$V_3(\text{H}_2\text{O}_2; \text{H}_2\text{O}) = V_3(\text{H}_2\text{O}) = V_1 \alpha k_2 x / f \quad (1)$$

$$= V_1 \alpha x / f_m \quad (1a)$$

In this equation⁷, V_1 is the initiation rate, $l = [\text{H}_2]$ and $x = [\text{O}_2]$ are the molar concentrations of the reactants with $l \gg x$; $\alpha = k_3/k_4$ is the ratio of the rate constants of the competing (parallel) reactions; $k_2 = \alpha \sqrt{2k_5 V_1} / x_m^2$ is the rate constant of reaction 2 of hydrogen atom addition to the oxygen molecule, whose analytical expression is

⁴The planar, six-atom, cyclic, hydrogen-bonded dimer $(\text{HO}_2^{\bullet})_2$ was calculated using quantum chemical methods (B3LYP density functional theory) [40]. The hydrogen bond energy is 47.7 and 49.4 kJ mol^{-1} at 298 K for the triplet and singlet states of the dimer, respectively.

⁵It is impossible to make a sharp distinction between the two-step bimolecular interaction of three species *via* the equilibrium formation of the labile intermediate O_4 and the elementary trimolecular reaction $\text{O}_2 + \text{H}^{\bullet} \rightarrow \text{HO}_4^{\bullet}$.

⁶For example, the ratio of the rate constants of the bimolecular disproportionation and dimerization of free radicals at room temperature is $k(\text{HO}^{\bullet} + \text{HO}_2^{\bullet})/[2k(2\text{HO}^{\bullet})2k(2\text{HO}_2^{\bullet})]^{0.5} = 2.8$ in the atmosphere [44] and $k(\text{H}^{\bullet} + \text{HO}^{\bullet})/[2k(2\text{H}^{\bullet})2k(2\text{HO}^{\bullet})]^{0.5} = 1.5$ in water [3]. These values that are fairly close to unity.

⁷This equation can be used to describe a wide range of nonbranched-chain reactions of addition of any free radicals to the C=C bonds of unsaturated hydrocarbons, alcohols, etc., which result in the formation of molecular 1:1 adducts in binary reaction systems of saturated and unsaturated components [1, 9, 10].

obtained by the solution to the quadratic equation derived from the condition of the extremum of the rate function $\partial V_3/\partial x = 0$; l_m and x_m are the concentrations of the components l and x , respectively, at the maximum of the function; $f = k_2x^2 + (\alpha l + x)\sqrt{2k_3V_1}$ and $f_m = x^2 + (\alpha l + x)x_m^2/\alpha l_m$; and $2k_5$ is the rate constant of hydrogen atom recombination reaction 5 considered to be bimolecular in this approximation.

The rate constant $2k_5$ in the case of the pulsed radiolysis of ammonia–oxygen (+ argon) gaseous mixtures at a total pressure of 10^5 Pa and a temperature of 349 K was calculated to be $1.6 \times 10^8 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ [6] (a similar value of this constant for the gas phase was reported in an earlier publication [48]). Pagsberg *et al.* [6] found that the dependence of the yield of the intermediate HO^\bullet on the oxygen concentration has a maximum close to $5 \times 10^{-4} \text{ mol dm}^{-3}$. In the computer simulation of the process, they considered the strongly exothermic reaction $\text{HO}_2^\bullet + \text{NH}_3 \rightarrow \text{H}_2\text{O} + \text{NHOH}$, which is similar to reaction 3 in *Scheme*, whereas the competing reaction 4 was not taken into account.

The ratio of the rates of the competing reactions is $V_3/V_4 = \alpha l/x$ and the chain length is $\nu = V_3/V_1$. Equation (1a) was obtained by substitution of the rate constant k_2 into Eq. (1) with its analytical expression (in order to reduce the number of unknown parameters that are to be measured directly). The optimum concentration x_m of oxygen, at which the rate of oxidation is maximum, can be calculated from Eq. (1a) or the analytical expression for k_2 if other parameters that appear in these expressions are known.

The rates of nonchain formation of molecular hydrogen, hydrogen peroxide, and water in reactions 5, 6, and 7 of quadratic chain termination are as follows:

$$V_5 = V_1^2 2k_5 (\alpha l + x)^2 / f^2, \quad (2)$$

$$2V_6 = 2V_1 \sqrt{2k_3V_1} (\alpha l + \beta + x) k_2 x^2 / f^2, \quad (3)$$

$$V_7 = V_1 (k_2 x^2) / f^2. \quad (4)$$

The designations of the parameters in this equation are the same as in Eq. (1).

The rate of the nonbranched-chain free-radical oxidation of hydrogen is a complex function of the rates of formation and disappearance of H^\bullet atoms and HO_4^\bullet radicals: $V_1 + V_{3,3'} - V_4 - V_5 + V_7$. Unlike the dependences of the rates V_4 ($V_4 \leq V_1$), V_5 and V_7 , the dependences of the rates V_2 , $V_{3,3'}$, and $2V_6$ on the oxygen concentration x show a maximum.

Equation (1) under the conditions (a) $k_2x^2 \ll (\alpha l + x)\sqrt{2k_3V_1}$, $\alpha l \gg x$ and (b) $k_2x^2 \gg (\alpha l + x)\sqrt{2k_3V_1}$, corresponding to the ascending and descending branches of the curve with a maximum, can be transformed into simple equations which allow preliminary to estimate the

parameters α and k_2 and express directly and inversely proportional functions of concentration x :

$$V_3 = \sqrt{V_1} k_2 x / \sqrt{2k_5}, \quad (5)$$

$$V_3 = V_1 \alpha l / \varphi x, \quad (6)$$

where $\varphi = 2$ at maximum when $k_2x^2 \cong (\alpha l + x)$ and $\varphi = 1$ at the decreasing part of the curve.

In the case of nonchain hydrogen oxidation *via* the above addition reaction ($\text{H}^\bullet + \text{O}_4 \xrightarrow{k_{add}} \text{HO}_4^\bullet$), the formation rates of the molecular oxidation products in reactions 6 and 7 (*Scheme*, $k_2 = k_3 = k_4 = 0$) are defined by modified Eqs. (3) and (4) in which $\beta = 0$, $(\alpha l + x)$ is replaced with 1, and k_2 is replaced with $k_{add}K_{eq}$ ($k_{add}K_{eq}$ is the effective rate constant of H^\bullet addition to the O_4 dimer, $K_{eq} = k/k'$ is the equilibrium constant of the reversible

reaction $2\text{O}_2 \xrightleftharpoons[k']{k} \text{O}_4$ with $k' \gg k_{add}[\text{H}^\bullet]$). The formation

rates of the stable products of nonchain oxidation ($k_3 = 0$), provided that either reactions 2 and 4 or reaction 2 alone ($k_4 = 0$) occurs (*Scheme*; in the latter case, reactions 6 and 7 involve the HO_2^\bullet radical rather than HO_4^\bullet), are given by modified Eqs. (3) and (4), in which $(\alpha l + x)$ replaced with 1, and x^2 replaced with x . In the latter case, the HO_2^\bullet radical, rather than HO_4^\bullet , takes part in reactions 6 and 7.

It is important to note that, if in the *Scheme* chain initiation *via* reaction 1 is due to the interaction between molecular hydrogen and molecular oxygen yielding the hydroxyl radical HO^\bullet instead of H^\bullet atoms and if this radical reacts with an oxygen molecule (reaction 4) to form the hydrotrioxyl radical HO_3^\bullet (which was obtained in the gas phase by neutralization reionization (NR) mass spectrometry [33] and has a lifetime of $>10^{-6}$ s at 298 K) and chain termination takes place *via* reactions 5–7 involving the HO^\bullet and HO_3^\bullet radicals instead of H^\bullet and HO_4^\bullet , respectively, the expression for the water chain formation rates (the chain formation of hydrogen peroxide in this kinetic model does not occur) derived in the same way will appear as a fraction rational function of the oxygen concentration x without a maximum: $V_3(\text{H}_2\text{O}) = V_3'(\text{H}_2\text{O}) = V_1 k_3 l / (k_4 x + \sqrt{2k_3V_1})$.

Curve 2 in Fig. 1 describes, in terms of the overall equation $V_{3,7} = V_1 x (\alpha l f_m + x^3) / f_m^2$ for the rates of reactions 3 and 7 (which was derived from Eqs. (1a) and (4), respectively, at that Eq. (4) in the form [49] of $V_7 = V_1 x^4 / f_m^2$ (4a) in which k_2 is replaced with its analytical expression $\alpha l_m \sqrt{2k_3V_1} / x_m^2$ derived from Eq. (1)), the dependence of the hydrogen peroxide formation rate (minus the rate $V_{\text{H}_2\text{O}_2} = 5.19 \times 10^{-8} \text{ mol dm}^{-3} \text{ s}^{-1}$ of the primary formation of hydrogen peroxide after completion of the reactions in spurs) on the concentration of dissolved oxygen during the γ -radiolysis of water saturated with hydrogen (at the initial concentration $7 \times$

10^{-4} mol dm $^{-3}$) at 296 K [7]. These data were calculated in the present work from the initial slopes of hydrogen peroxide buildup versus dose curves for a ^{60}Co γ -radiation dose rate of $P = 0.67$ Gy s $^{-1}$ and absorbed doses of $D \cong 22.5\text{--}304.0$ Gy. The following values of the primary radiation-chemical yield G (species per 100 eV of energy absorbed) for water γ -radiolysis products in the bulk of solution at pH 4–9 and room temperature were used (taking into account that $V = GP$ and $V_1 = G_{\text{H}}P$): $G_{\text{H}_2\text{O}_2} = 0.75$ and $G_{\text{H}} = 0.6$ (initiation yield) [3]; $V_1 = 4.15 \times 10^{-8}$ mol dm $^{-3}$ s $^{-1}$; $2k_5 = 2.0 \times 10^{10}$ dm 3 mol $^{-1}$ s $^{-1}$ [3]. As can be seen from Fig. 1, the best description of the data with an increase in the oxygen concentration in water is attained when the rate V_7 of the formation of hydrogen peroxide *via* the nonchain mechanism in the chain termination reaction 7 (curve 1, $\alpha = (8.5 \pm 2) \times 10^{-2}$) is taken into account in addition to the rate V_3 of the chain formation of this product *via* the propagation reaction 3 (dashed curve 2, $\alpha = 0.11 \pm 0.026$). The rate constant of addition reaction 2 determined from α is substantially underestimated: $k_2 = 1.34 \times 10^7$ (vs. 2.0×10^{10} [3]) dm 3 mol $^{-1}$ s $^{-1}$. The difference can be due to the fact that the radiation-chemical specifics of the process were not considered in the kinetic description of the experimental data. These include oxygen consumption *via* reactions that are not involved in the hydrogen oxidation *Scheme* and reverse reactions resulting in the decomposition of hydrogen peroxide by intermediate products of water radiolysis (e_{aq}^- , H^\cdot , HO^\cdot), with the major role played by the hydrated electron [3].

III. CONCLUSIONS

Thus, the addition reaction of the HO_2^\cdot radical that possesses an elevated energy in statu nascendi with the oxygen molecule (at sufficiently high oxygen concentrations) to give the HO_4^\cdot radical was used for the first time in the kinetic description of the initiated hydrogen oxidation at moderate temperature and pressure [9, 10]. This reaction is endothermic and competes with the chain propagation reaction *via* the H^\cdot atom. The HO_4^\cdot radical generated in the former reaction has a low reactivity and inhibits the chain reaction.⁸

The above data concerning the competition kinetics of the nonbranched-chain addition of hydrogen atoms to the multiple bonds of the oxygen molecules make it possible to describe, using rate equations (1a) and (4a), obtained by quasi-steady-state treatment, the peaking experimental dependences of the formation rates of molecular 1:1 adduct H_2O_2 on the concentration of the oxygen over the entire range of its variation in binary system (Fig. 1). In such reaction systems consisting of saturated and

unsaturated components [51, 52], the unsaturated compound (in this case the O_2) is both a reactant and an autoinhibitor, specifically, a source of low-reactive free radicals (in this case the HO_4^\cdot radicals) shortening kinetic chains.

The progressive inhibition of the nonbranched-chain processes, which takes place as the concentration of the unsaturated compound is raised (after the maximum process rate is reached), can be an element of the self-regulation of the natural processes that returns them to the stable steady state.

Using mechanism of the nonbranched-chain free-radical hydrogen oxidation considered here, it has been demonstrated that, in the Earth's upper atmosphere, the decomposition of O_3 in its reaction with the HO^\cdot radical can occur *via* the addition of the latter to the ozone molecule, yielding the HO_4^\cdot radical, which is capable of efficiently absorbing UV radiation [32].

REFERENCES

- [1] M. M. Silaev and L. T. Bugaenko, "Mathematical Simulation of the Kinetics of Radiation Induced Hydroxyalkylation of Aliphatic Saturated Alcohols", *Radiation Physics and Chemistry*, 1992, vol. 40, no. 1, pp. 1–10.
- [2] M. M. Silaev, "Applied Aspects of the γ -Radiolysis of C $_1$ –C $_4$ Alcohols and Binary Mixtures on Their Basis", *Khimiya Vysokikh Energii*, Vol. 36, No. 2, 2002, pp. 97–101, English Translation in: *High Energy Chemistry*, 2002, vol. 36, no. 2, pp. 70–74.
- [3] A. K. Pikaev, "Sovremennaya radiatsionnaya khimiya. Radioliz gazov i zhidkostei" ("Modern Radiation Chemistry: Radiolysis of Gases and Liquids"), Nauka, Moscow, 1986.
- [4] H. A. Smith and A. Napravnik, "Photochemical Oxidation of Hydrogen", *Journal of the American Chemistry Society*, 1940, vol. 62, no. 1, pp. 385–393.
- [5] E. J. Badin, "The Reaction between Atomic Hydrogen and Molecular Oxygen at Low Pressures. Surface Effects", *Journal of the American Chemistry Society*, 1948, vol. 70, no. 11, pp. 3651–3655.
- [6] P. B. Pagsberg, J. Eriksen, and H. C. Christensen, "Pulse Radiolysis of Gaseous Ammonia–Oxygen Mixtures", *The Journal of Physical Chemistry*, 1979, vol. 83, no. 5, pp. 582–590.
- [7] N. F. Barr and A. O. Allen, "Hydrogen Atoms in the Radiolysis of Water", *The Journal of Physical Chemistry*, 1959, vol. 63, no. 6, pp. 928–931.
- [8] M. M. Silaev, "A New Competitive Kinetic Model of Radical Chain Oxidation: Oxygen as an Autoinhibitor", *Biofizika*, 2001, vol. 46, no. 2, pp. 203–209, English Translation in: *Biophysics*, 2001, vol. 46, no. 2, pp. 202–207.
- [9] M. M. Silaev, "The Competition Kinetics of Nonbranched Chain Processes of Free-Radical Addition to Double Bonds of Molecules with the Formation of 1:1 Adducts and the Inhibition by the Substrate", *Oxidation Communication*, 1999, vol. 22, no. 2, pp. 159–170.
- [10] M. M. Silaev, "The Competition Kinetics of Radical-Chain Addition", *Zhurnal Fizicheskoi Khimii*, Vol. 73, No. 7, 1999, pp. 1180–1184, English Translation in:

⁸Note that in the case of similar nonbranched-chain (i.e., "slow") oxidation of RH hydrocarbons, the corresponding reactive RO_2^\cdot and low-reactive RO_4^\cdot radicals participate in the process [50]. The only difference between the kinetic model of oxidation and the kinetic model of the chain addition of 1-hydroxyalkyl radicals to the free (unsolvated) form of formaldehyde in nonmethanolic alcohol–formaldehyde systems [1, 9, 10] is that in the former does not include the formation of the molecular 1:1 adduct *via* reaction 4.

- Russian Journal of Physical Chemistry*, 1999, vol. 73, no. 7, pp. 1050–1054.
- [11] M. M. Silaev, “Competitive Mechanism of the Nonbranched Radical Chain Oxidation of Hydrogen Involving the Free Cyclohydroperoxy Radical $[\text{OO}\cdots\text{H}\cdots\text{OO}]^*$, which Inhibits the Chain Process”, *Khimiya Vysokikh Energii*, 2003, vol. 37, no. 1, pp. 27–32, English Translation in: *High Energy Chemistry*, 2003, vol. 37, no. 1, pp. 24–28.
- [12] M. M. Silaev, “Simulation of the Initiated Addition of Hydrocarbon Free Radicals and Hydrogen Atoms to Oxygen via a Nonbranched Chain Mechanism”, *Teoreticheskie Osnovy Khimicheskoi Tekhnologii*, 2007, vol. 41, no. 6, pp. 634–642, English Translation in: *Theoretical Foundation of Chemical Engineering*, 2007, vol. 41, no. 6, pp. 831–838.
- [13] M. M. Silaev, “Simulation of Initiated Nonbranched Chain Oxidation of Hydrogen: Oxygen as an Autoinhibitor”, *Khimiya Vysokikh Energii*, 2008, vol. 42, no. 2, pp. 124–129, English Translation in: *High Energy Chemistry*, 2008, vol. 42, no. 1, pp. 95–100.
- [14] N. N. Semenov, “Tsepnye reaktsii” (“Chain Reactions”), Goskhimtekhnizdat, Leningrad, 1934, pp. 241, 203.
- [15] J. S. Francisco and I. H. Williams, “The Thermochemistry of Polyoxides and Polyoxy Radicals”, *International Journal of Chemical Kinetics*, 1988, vol. 20, no. 6, pp. 455–466.
- [16] D. J. McKay and J. S. Wright, “How Long Can You Make an Oxygen Chain?”, *Journal of the American Chemistry Society*, 1998, vol. 120, no. 5, pp. 1003–1013.
- [17] N. P. Lipikhin, “Dimers, Clusters, and Cluster Ions of Oxygen in the Gas Phase”, *Uspekhi Khimii*, 1975, vol. 44, no. 8, pp. 366–376.
- [18] S. D. Razumovskii, “Kislorod – elementarnye formy i svoistva” (“Oxygen: Elementary Forms and Properties”), *Khimiya*, Moscow, 1979.
- [19] K. M. Dunn, G. E. Scuseria, and H. F. Schaefer III, “The infrared spectrum of cyclotetraoxygen, O_4 : a theoretical investigation employing the single and double excitation coupled cluster method”, *The Journal of Chemical Physics*, 1990, vol. 92, no. 10, pp. 6077–6080.
- [20] L. Brown and V. Vaida, “Photoreactivity of Oxygen Dimers in the Ultraviolet”, *The Journal of Physical Chemistry*, 1996, vol. 100, no. 19, pp. 7849–7853.
- [21] V. Aquilanti, D. Ascenzi, M. Bartolomei, D. Cappelletti, S. Cavalli, M. de Castro-Vitores, and F. Pirani, “Molecular Beam Scattering of Aligned Oxygen Molecules. The Nature of the Bond in the $\text{O}_2\text{--O}_2$ Dimer”, *Journal of the American Chemistry Society*, 1999, vol. 121, no. 46, pp. 10794–1080.
- [22] F. Cacace, G. de Petris, and A. Troiani, “Experimental Detection of Tetraoxygen”, *Angewandte Chemie, International Edition* (in English), 2001, vol. 40, no. 21, pp. 4062–4065.
- [23] H. S. Taylor, “Photosensitisation and the Mechanism of Chemical Reactions”, *Transactions of the Faraday Society*, 1926, vol. 21, no. 63(3), pp. 560–568.
- [24] S. N. Foner and R. L. Hudson, “Mass spectrometry of the HO_2 free radical”, *The Journal of Chemical Physics*, 1962, vol. 36, no. 10, p. 2681.
- [25] S. N. Foner and R. L. Hudson, “Mass spectrometry of the HO_2 free radical”, *The Journal of Chemical Physics*, 1962, vol. 36, no. 10, p. 2681.
- [26] P. D. Lightfoot, B. Veyret, and R. Lesclaux, “Flash Photolysis Study of the $\text{CH}_3\text{O}_2 + \text{HO}_2$ Reaction between 248 and 573 K”, *The Journal of Physical Chemistry*, 1990, vol. 94, no. 2, pp. 708–714.
- [27] P. Smith, “The HO_3 and HO_4 Free Radicals”, *Chemistry and Industry*, 1954, no. 42, pp. 1299–1300.
- [28] A. J. B. Robertson, “A Mass Spectral Search for H_2O_4 and HO_4 in a Gaseous Mixture Containing HO_2 and O_2 ”, *Chemistry and Industry*, 1954, no. 48, p. 1485.
- [29] D. Bahnemann and E. J. Hart, “Rate Constants of the Reaction of the Hydrated Electron and Hydroxyl Radical with Ozone in Aqueous Solution”, *The Journal of Physical Chemistry*, 1982, vol. 86, no. 2, pp. 252–255.
- [30] G. C. Pimentel and A. L. McClellan, “The Hydrogen Bond”, L. Pauling, Editor, Freeman, San Francisco, 1960, p. 200.
- [31] “Vodorodnaya svyaz’: Sbornik statei” (“The Hydrogen Bonding: Collection of Articles”) N. D. Sokolov, Editor, Nauka, Moscow, 1981.
- [32] J. Staehelin, R. E. Bühler, and J. Hoigné, “Ozone Decomposition in Water Studied by Pulse Radiolysis. 2. OH and HO_4 as Chain Intermediates”, *The Journal of Physical Chemistry*, 1984, vol. 88, no. 24, pp. 5999–6004.
- [33] F. Cacace, G. de Petris, F. Pepi, and A. Troiani, “Experimental Detection of Hydrogen Trioxide”, *Science*, 1999, vol. 285, no. 5424, pp. 81–82.
- [34] R. F. Bühler, J. Staehelin, and J. Hoigné, “Ozone Decomposition in Water Studied by Pulse Radiolysis. 1. HO_2/O_2^- and HO_3/O_3^- as Intermediates”, *The Journal of Physical Chemistry*, 1984, vol. 88, no. 12, pp. 2560–2564.
- [35] I. V. Trushkov, M. M. Silaev, and N. D. Chuvylkin, “Acyclic and Cyclic Forms of the Radicals HO_4^* , CH_3O_4^* , and $\text{C}_2\text{H}_5\text{O}_4^*$: Ab Initio Quantum Chemical Calculations”, *Izvestiya Akademii Nauk, Ser.: Khimiya*, 2009, no. 3, pp. 479–482, English Translation in: *Russian Chemical Bulletin, International Edition*, 2009, vol. 58, no. 3, pp. 489–492.
- [36] A. Mansergas, J. M. Anglada, S. Olivella, M. F. Ruiz-López, “On the Nature of the Unusually Long OO Bond in HO_3 and HO_4 Radicals”, *Phys. Chem. Chem. Phys.*, 2007, vol. 9, no. 44, pp. 5865–5873.
- [37] W. Wong and D. D. Davis, “A Flash Photolysis Resonance Fluorescence Study of the Reactions of Atomic Hydrogen and Molecular Oxygen: $\text{H} + \text{O}_2 + \text{M} \rightarrow \text{HO}_2 + \text{M}$ ”, *International Journal of Chemical Kinetics*, 1974, vol. 6, no. 3, pp. 401–416.

Free-Radical Nonbranched-Chain Oxidation of Hydrogen

- [38] S. W. Benson, "Thermochemical Kinetics: Methods for the Estimation of Thermochemical Data and Rate Parameters", 2nd Edition, Wiley, New York, 1976.
- [39] Yagodovskaya, T.V. and Nekrasov, L.I., "Use of Infrared Spectroscopy to Study Frozen Hydrogen–Oxygen Systems Containing Hydrogen Polyoxides", *Zhurnal Fizicheskoi Khimii*, 1977, vol. 51, no. 10, pp. 2434–2445.
- [40] X. Xu, R. P. Muller, and W. A. Goddard III, "The Gas Phase Reaction of Singlet Dioxygen with Water: a Water-Catalyzed Mechanism", *Proceedings of the National Academy Sciences of the United States of America*, 2002, vol. 99, no. 6, pp. 3376–3381.
- [41] E. T. Seidl and H. F. Schaefer III, "Is There a Transition State for the Unimolecular Dissociation of Cyclotetraoxygen (O₄)?", *The Journal of Chemical Physics*, 1992, vol. 96, no. 2, pp. 1176–1182.
- [42] R. Hernández-Lamonedá and A. Ramírez-Solís, "Reactivity and Electronic States of O₄ along Minimum Energy Paths", *The Journal of Chemical Physics*, 2000, vol. 113, no. 10, pp. 4139–4145.
- [43] A. J. C. Varandas and L. Zhang, "Test Studies on the Potential Energy Surface and Rate Constant for the OH + O₃ Atmospheric Reaction", *Chemical Physics Letters*, 2000, vol. 331, nos. 5–6, pp. 474–482.
- [44] "Atmosfera. Spravochnik" ("Atmosphere: A Handbook"), Gidrometeoizdat, Leningrad, 1991.
- [45] H. Okabe, "Photochemistry of Small Molecules", Wiley, New York, 1978.
- [46] A. B. Nalbandyan and V. V. Voevodskii, "Mekhanizm okisleniya i gorenija vodoroda" ("Mechanism of Hydrogen Oxidation and Combustion"), V. N. Kondrat'ev, Editor, Akad. Nauk SSSR, Moscow, 1949.
- [47] L. Bateman, "Olefin Oxidation", *Quarterly Reviews*, 1954, vol. 8, no. 2, pp. 147–167.
- [48] A. W. Boyd, C. Willis, and O. A. Miller, "A Re-examination of the Yields in the High Dose Rate Radiolysis of Gaseous Ammonia", *Canadian Journal of Chemistry*, 1971, vol. 49, no. 13, pp. 2283–2289.
- [49] M. M. Silaev, "Competition Mechanism of Substre-Inhibited Radical Chain Addition to Double Bond", *Neftekhimiya*, 2000, vol. 40, no. 1, pp. 33–40, English Translation in: *Petroleum Chemistry*, 2000, vol. 40, no. 1, pp. 29–35.
- [50] M. Silaev, Mathematical Simulation of Kinetics of Radiation-Induced Free-Radical Oxidation by Nonbranched-Chain Mechanism // *Recent Innovations in Chemical Engineering*, 2015, vol. 8, no. 2, pp. 112–124.
- [51] M. M. Silaev, "Low-reactive Free Radicals Inhibiting Nonbranched Chain Processes of Addition", *Biofizika*, 2005, vol. 50, no. 4, pp. 585–600, English Translation in: *Biophysics*, 2005, vol. 50, no. 4, pp. 511–524.
- [52] M. M. Silaev, Kinetics of the Free-Radical Nonbranched-Chain Addition // *American Journal of Polymer Science and Technology*, 2017, vol. 3, no. 3, pp. 29–49.