

Models in Country Scale Carbon Accounting of Forest Soils

Mikko Peltoniemi, Esther Thürig, Stephen Ogle, Taru Palosuo, Marion Schrumpf, Thomas Wutzler, Klaus Butterbach-Bahl, Oleg Chertov, Alexander Komarov, Alexey Mikhailov, Annemieke Gärdenäs, Charles Perry, Jari Liski, Pete Smith and Raisa Mäkipää

Peltoniemi, M., Thürig, E., Ogle, S., Palosuo, T., Schrumpf, M., Wutzler, T., Butterbach-Bahl, K., Chertov, O., Komarov, A., Mikhailov, A., Gärdenäs, A., Perry, C., Liski, J., Smith, P. & Mäkipää, R. 2007. Models in country scale carbon accounting of forest soils. *Silva Fennica* 41(3): xxx–xxx.

Countries need to assess changes in the carbon stocks of forest soils as a part of national greenhouse gas (GHG) inventories under the United Nations Framework Convention on Climate Change (UNFCCC) and the Kyoto Protocol (KP). Since measuring these changes is expensive, it is likely that many countries will use alternative methods to prepare these estimates. We reviewed seven well-known soil carbon models from the point of view of preparing country-scale soil C change estimates. We first introduced the models and explained how they incorporated the most important input variables. Second, we evaluated their applicability at regional scale considering commonly available data sources. Third, we compiled references to data that exist for evaluation of model performance in forest soils. A range of process-based soil carbon models differing in input data requirements exist, allowing some flexibility to forest soil C accounting. Simple models may be the only reasonable option to estimate soil C changes if available resources are limited. More complex models may be used as integral parts of sophisticated inventories assimilating several data sources. Currently, measurement data for model evaluation are common for agricultural soils, but less data have been collected in forest soils. Definitions of model and measured soil pools often differ, ancillary model inputs require scaling of data, and soil C measurements are uncertain. These issues complicate the preparation of model estimates and their evaluation with empirical data, at large scale. Assessment of uncertainties that accounts for the effect of model choice is important part of inventories estimating large-scale soil C changes. Joint development of models and large-scale soil measurement campaigns could reduce the inconsistencies between models and empirical data, and eventually also the uncertainties of model predictions.

Keywords decomposition, greenhouse gas inventory, IPCC, national forest inventory, regional and national modeling, soil carbon, soil model

Addresses (coresponding) *Peltoniemi* Finnish Forest Research Institute, Vantaa, Finland

E-mail mikko.peltoniemi@metla.fi (Other authors' addresses listed on the next page)

Received 25 August 2006 **Revised** 23 April 2007 **Accepted** 4 July 2007

Available at <http://www.metla.fi/silvafennica/full/sf41/sf413xxx.pdf>

1 Introduction

According to the UN Framework Convention on Climate Change and the Kyoto Protocol, countries need to include changes in soil and litter carbon pools in their annual greenhouse gas inventory submissions (UNFCCC 1992, UNFCCC 1997). Reliable methods to estimate soil carbon stock changes are needed for reporting, and for the assessment of soil response to changing climate.

The Intergovernmental Panel on Climate Change provides guidelines for conducting these inventories and has ranked approaches from Tier 1 to Tier 3 (IPCC 2003). The Tier 1 approach is the simplest for estimating changes and soil C stocks, requiring only country-specific data on forest land use. For forest lands, this method assumes no change in soil C stocks due to management. If there are changes in soil C stocks associated with other activities in forest lands than only land use, either a Tier 2 or Tier 3 approach is needed. Tier 2 also uses simple empirical relationships that are derived from country-specific data. Tier 3 approaches are completely country-specific, and presumably the most accurate.

The most straightforward way of preparing Tier 3 estimates would be repeated sampling of changes in soil C. However, presently, only a few countries, the UK (Bellamy et al. 2005), Belgium (Lettens et al. 2005a), and Sweden (Ståhl et al. 2004), have soil surveys suitable to provide nationwide estimates of soil C stock changes. Establishing such surveys in all countries is unrealistic because of high costs and extensive effort needed to collect the data for nationwide sampling. Costs increase greatly with increasing variation in soil properties, as is often the case in forest soils.

Flux-based measurements of ecosystems provide information on changes in carbon stocks as a whole (see e.g. Baldocchi 2003), but partitioning fluxes into vegetation and soil components requires either modelling of soil carbon dynamics or measuring plant detritus inputs and heterotrophic soil respiration. The present grid of measurement stations is also too sparse for reliable national estimates with flux measurements alone. Regional scaling has been made

with satellite measured photosynthetically active radiation (PAR) and models of net ecosystem exchange (NEE), and in conjunction with measurements made from research aircraft (Miglietta et al. 2007).

Changes in soil C stocks can also be estimated across a country by combining input data with process-based models (Ogle and Paustian 2005). This approach is another example of a Tier 3 method, and is the focus of this review paper. The details of this approach vary. For example, Liski et al. (2006) presented a calculation based on aggregated forest inventory data and a dynamic soil model. Post et al. (2001) proposed a combination of process-based models and multiple data sources such as eddy-covariance measurements, forest inventory data, and soil measurements. Lagergren et al. (2006) calibrated ecosystem model Biome-BGC with eddy covariance measurements, and used it in conjunction with forest inventory data to estimate the carbon balance of Swedish forests. Several countries have chosen, and are already using, either a fully or partly process-based modelling approach for the nationwide reporting of changes of carbon in forest soils (UNFCCC 2006).

Authors' addresses *Peltoniemi & Mäkipää*: Finnish Forest Research Institute, Vantaa, Finland; *Thürig*: Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland; *Palosuo & Thürig*: European Forest Institute, Joensuu, Finland; *Ogle*: Natural Resources Ecology Laboratory, Colorado State University, Fort Collins, USA; *Schrumpf & Wutzler*: Max-Planck-Institute for Biogeochemistry, Jena, Germany; *Butterbach-Bahl*: Institute for Meteorology and Climate Research, Forschungszentrum Karlsruhe GmbH, Garmisch-Partenkirchen, Germany; *Chertov*: St. Petersburg State University, St. Petersburg-Peterhof, Russia; *Komarov & Mikhailov*: Institute of Physicochemical and Biological Problems in Soil Science of Russian Academy of Sciences, Pushchino, Russia; *Gårdenäs*: Dept. of Soil Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden; *Perry*: USDA Forest Service, Northern Research Station, St. Paul, MN USA; *Liski*: Finnish Environment Institute, Helsinki, Finland; *Smith*: School of Biological Sciences, University of Aberdeen, Aberdeen, UK

A few comprehensive soil model reviews have been published in recent years. McGill (1996) assessed the structure of 10 soil models, and found that the trend was toward kinetic compartmentalization of SOM. Smith et al. (1997) compared performance of nine SOM models against repeated long-term measurement from agricultural, grassland and woodland soils, including several models discussed in this review. Rodrigo et al. (1997) compared temperature and moisture functions of selected models, but only one of those models is included in this study (*SOILN*). Izaurre et al. (2001) was the only review that compared models for the purpose of selecting the best performing model for a regional application.

All previous model comparisons were done for land-uses other than forests except for two woodland / forestry sites used in Smith et al. (1997). Therefore, our goal was to review existing soil models that are available for the estimation of carbon stock changes in forest soils, providing an overview to scientists and officials involved with national GHG inventories.

The objective of this study is to analyze how different soil carbon models can be applied to estimate short-term changes in soil carbon over large spatial scales. Specifically, (1) we review 7 soil models used for estimating forest soil C stock changes (*Century*, *Forest-DNDC*, *ROMUL*, *RothC*, *SOILN*, *Yasso*, and a statistical model *Forcarb*), (2) we evaluate the availability of input data for these models, and (3) compile references to relevant datasets that could be used for their evaluation.

2 Models Estimating Soil Carbon Fluxes

2.1 Decomposition Process

Forest and agricultural soils differ in many respects; they experience different management/disturbance regime, and there are differences in their vegetation and biota. Some authors have suggested that different model structures might be needed for forest and agricultural soils (Li et al. 2000, Chertov et al. 2001) but many models have been applied to both types of soil, although

some with modified parameterization (Peng et al. 1998, Eckersten et al. 1999, Kätterer et al. 1999, Falloon and Smith 2002, Falloon et al. 2002, Peng et al. 2002, Smith et al. 2006).

Decomposition is mediated mainly by the activity of soil microbes, fungi and fauna, but their specific population dynamics and explicit contribution to decomposition is rarely described in soil models (McGill 1996). A few models provide a very coarse scale description of biological control: *ROMUL* regulates the translocation of SOM between soil layers with soil biota (Chertov et al. 2001), and decomposition in *SOILN* (Eckersten and Beier 1998) and *Q-model* (Rolff and Ågren 1999, Ågren and Hyvönen 2003) is controlled by microbial biomass. Most models assume that the size of the microbial pool does not explicitly restrict decomposition, but rather that decomposition is limited by variables known to be correlated with microbial activity. Smith (2001, 2002) reviews the representation of decomposition processes in different SOM models.

In most models, microbial activity is expressed in the decomposition rates of model pools, which are typically first-order rate constants regulated by variables describing ambient conditions and properties of the soil matrix. Compounds belonging to more stable fractions of SOM require higher activation energies to decompose (Davidson and Janssens 2006). The complexity of degrading compounds creates a continuum of activation energies, which is usually approximated with several pools differing in turnover time. Empirical data and modelling studies also imply that the temperature effect on decomposition cannot be adequately approximated with only one soil pool (Kätterer et al. 1998, Davidson et al. 2000, Knorr et al. 2005, Davidson and Janssens 2006). All of the models included in this review describe decomposition of SOM with a multi-pool approach (Table 3).

Besides temperature, decomposition of litter or SOM can be affected by several variables, such as topography (mostly via moisture and temperature), parent material and management (Jenny 1941), litter quality, nitrogen or other macronutrients (Melillo et al. 1982, Prescott 1995, Berg 2000), heavy metals (Berg and McLaugherty 2003), and chemical weathering (Sverdrup 1990, Sverdrup et al. 1995). SOM decomposition may

also be influenced by chemical or physical protection through occlusion of SOM in complexes with clay mineral or encapsulation within soil aggregates (Oades 1988, Christensen 1996, Elliot et al. 1996, Six et al. 2002), as well as influenced by drought, flooding or freeze/thaw cycles (Davidson and Janssens 2006).

The complexity of the decomposition process and large uncertainties in empirical data make it difficult to develop a completely accurate model as well as to parameterize exceedingly sophisticated models.

2.2 Process-based Models - Key Factors Affecting Decomposition

Below we describe how the most typical input variables (i.e. temperature, moisture, soil texture and nitrogen) influence the decomposition process as represented in six common models, including *SOILN* (Eckersten and Beier 1998), *RothC* (Coleman and Jenkinson 1996), *ROMUL* (Chertov et al. 2001, Bykhovets and Komarov 2002), *Century* (Parton et al. 1987, Parton et al. 1994), *Yasso* (Liski et al. 2005) and *Forest-DNDC* (Li et al. 2000, Stange et al. 2000). The minimum required input information needed by the model users is presented in Table 1, and the model-specific influences on the decomposition process are described in Table 2. Model initialization is also discussed due to its importance for estimating accurate trends.

2.2.1 Temperature

In all models, temperature affects a rate modifier that multiplies the decomposition rates of one or several compartments. Main differences among the models arise from use of air or soil temperature as the input data, exact formulation of the temperature models, and their time steps.

In *SOILN*, daily soil temperature of each soil layer is simulated with the water and heat model *CoupModel* (Jansson and Karlberg 2002) using information of daily air temperature and global radiation. The simulated soil temperatures are used as driving variables to calculate decomposition rates. In *SOILN* there are three alternative

functions to describe the effect of soil temperature on decomposition, i) the Q_{10} , ii) the Q_{10} after threshold, and iii) the Ratkowsky functions (Ratkowsky et al. 1982). The Q_{10} function, originally introduced by van 't Hoff and Arrhenius in late 19th Century, implies that for each 10 degree increase in temperature, decomposition rates increase with the Q_{10} factor (Lloyd and Taylor 1994). The Q_{10} after threshold means that the Q_{10} relationship is applied at temperatures higher than a certain threshold, commonly 0 to 5 °C.

Forest-DNDC uses daily minimum and maximum air temperature as a model driver, and implements an O'Neill response function to describe the effect of temperature on decomposition (Stange 2007). This function yields an exponential increase in decomposition at lower temperatures ranges, but it has a distinct temperature optimum and a sharp decrease at higher temperatures. The effects of moisture and temperature on decomposition are combined by multiplication in many C-decomposition models response functions, including *Forest-DNDC* (see e.g. review by Rodrigo et al. 1997). Moreover, in the *Forest-DNDC* model, the advantage of using a multiplier for moisture and temperature effects on decomposition (i.e., continuity) is combined in a framework to evaluate the most limiting factor for microbial activity (Liebig's law). This was achieved by adding reciprocal values of the respective response functions for moisture and temperature. Using this approach, the factor most limiting for decomposition (temperature or soil moisture) dominates the relationship (Li et al. 2000, Stange et al. 2000).

In *RothC* (Coleman and Jenkinson 1996), mean monthly air temperature (°C) is used. Temperature effects are represented as a multiplicative rate modifier on the decomposition of active compartments. The relationship is close to linear between temperatures 10 and 35°C.

In *ROMUL* (Chertov et al. 2001), forest floor and soil temperatures are used to calculate non-linear rate modifiers for decomposition rates of aboveground and belowground litter portions correspondingly. These soil temperatures may be simulated by the *SCLISS* generator (Bykhovets and Komarov 2002), which uses monthly air temperature and soil texture parameters as input information. Temperature corrections that depend

Table 1. Minimum required input data according to variables and temporal resolution.

	<i>Yasso</i>	<i>ROMUL</i>	<i>SOILN</i>	<i>RothC</i>	<i>Forest-DNDC</i>	<i>CENTURY</i>
Time-step	Year	Month, other time-steps also possible	Day	Month	Day, part of the routines hourly or even sub-hourly	Month (daily version exists)
Litter input data	Amount of non-woody, fine woody and coarse woody litter	Amount of leaves, branches, stem, coarse and fine roots	C and N content of fine litter, coarse litter, humus and optionally in microbes	Amount of decomposable and resistant plant material	Plant material transferred during senescence to above and below-ground structural and metabolic litter pools	Plant material transferred during senescence to above and belowground structural and metabolic litter pools
Temperature (T) input	Annual average air T or effective T sum, monthly mean T (MMT) to calculate PET (May-Sep)	Mean monthly soil T in organic layer and in mineral layer or mean monthly air T as simulated with SCLISS	Daily soil temperature of each layer as simulated with <i>CoupModel</i>	Mean monthly air T	Daily min, max or alternatively average daily temperature	Monthly min and max T, (canopy biomass to calculate soil T)
Moisture input	Monthly precipitation (PPT) May-Sep	Monthly PPT and hydrological soil parameters (Bykhovets and Komarov 2002)	Soil moisture content, and soil water flows into and from the different soil layers (i.e. also infiltration and percolation flow). AET/PET as simulated with <i>CoupModel</i>	Monthly PPT and PET or open pan evaporation (Smith et al. 2005b, Smith et al. 2006)	Daily PPT, in case of the wetland model (Cui et al. 2005)	Monthly PPT from rainfall and snow, as well as irrigation inputs of water and losses from evapotranspiration, runoff and ground-water flow.
Texture input	–	Clay content (ROMUL). Loam and sand (SCLISS soil weather generator)	Hydraulic properties either measured or from a database (Jansson and Karlberg 2002)	Clay content	Clay content	Sand, silt and clay content
N input	–	N deposition data	N deposition and fertilization, initial N content of different plant parts	–	N deposition and fertilization	N deposition and fertilization, organic amendment N inputs
Nutrients (excl. N)	–	–	–	–	–	P,S
Initialization	Spin-up with steady state assumption or by allocating measured values with some assumptions	Measured or compiled data on SOM and N pools in organic layer and mineral topsoil	Measured C and N content of a mature forest in combination with steady state assumption	Either initialised to measured SOC (Smith et al. 2005b, Smith et al. 2006) or run to equilibrium using estimated inputs (Thürig et al. 2006)	Can be initialised with measured SOC values (Li et al. 2000, Kesik et al. 2005)	Typically initialized with spin-up (several 1000 years) under native vegetation, measured soil characteristics and mean climate for the site.

Table 2. Effect of environmental factors on modeled decomposition.

	<i>Yasso</i>	<i>ROMUL</i>	<i>SOILN</i>	<i>RothC</i>	<i>Forest-DNDC</i>	<i>CENTURY</i>
Effect of T and moisture on modelled decomposition (D)	D-rates adjusted by empirical regression models based on litter bag experiment (Liski et al. 2003)	Empirical regression models based on laboratory data (Chertov et al. 2001)	D-rates calculated using empirical functions and simulated values of daily soil temperature and moisture content	Soil moisture deficit and T affects rate modifier that multiplies the D-rates of active compartments. (Coleman and Jenkinson 1996)	Combined moisture and temperature factor used to modify potential D, nitrification or denitrification rates. (Li et al. 2000, Stange et al. 2000)	Soil moisture and temperature effects on D modeled with DEFAC relationship (multiplier combining temperature and moisture effects)
Effects of texture on modeled decomposition	–	Affects stable humus mineralization rate and soil moisture in SCLISS	Affects decomposition rate, N leaching, denitrification	Affects SOM stabilisation and soil water retention (Coleman and Jenkinson 1996)	Affects aeration, ammonium absorption, water movement and soil moisture, decomposition etc.	Affects moisture availability (Metherell et al. 1993) and decomposition rates (Parton et al. 1987, Parton et al. 1994).
Effect of N on modeled decomposition	–	N dynamics calculated in close interaction with SOM (soil C) dynamics	Dynamically coupled C and N model	–	Explicit N model, algorithm partly derived from DNDC (Li et al. 1992, Li et al. 2000)	Dynamically coupled C and N model.
Nutrients (excl. N)	–	–	–	–	–	P,S optional sub-models

on forest type are used in some cases due to difference between standard meteorological data and the temperature of forest floor and soil.

In *Century*, soil temperature is modeled as a function of daily maximum and minimum air temperatures, along with the influence of canopy cover on the radiation budget (Metherell et al. 1993). Temperature effects on decomposition vary across a range of temperatures according to empirically-derived relationships from decomposition studies (Parton et al. 1987). The influence of temperature on decomposition is combined with moisture into a single multiplicative factor (Parton et al. 1994).

In *Yasso* (Liski et al. 2005), the effect of temperature on decomposition is described with linear regression models that modify decomposition rates (Liski et al. 2003). The description of the effect of climate on decomposition in *Yasso* is based on geographically derived litterbag data. These regressions are scaled with decomposi-

tion rates at the reference locations to estimate rate modifiers for decomposition in each model compartment. Temperature variables used in the regression models are mean annual temperature (MAT) or effective temperature sum (DD0).

2.2.2 Soil Moisture

The effect of soil moisture on decomposition in the models is closely related or similar to the effect of temperature on decomposition.

In *SOILN*, daily soil moisture content of each soil layer is simulated with the water and heat model *CoupModel* (Jansson and Karlberg 2002) and they are used as driving variables to calculate decomposition rates. *CoupModel* includes the main processes influencing soil water balance for high latitudes such as precipitation (rain, snow), evaporation (transpiration, interception and soil evaporation) and losses like surface run-off,

percolation and drainage. Decomposition rates are assumed to be optimal for a defined soil moisture range. The user defines this range by two threshold values. The first threshold value is around the wilting point (i.e., minimum water content for optimal decomposition rate), while the second threshold is close to saturation (i.e., maximum water content for optimal decomposition rate). The relationship used to estimate the range of limiting water contents can be described with a linear, concave or convex function. The user may also define specific relationships for different processes, such as decomposition and nitrification.

Forest-DNDC uses total daily rainfall as input to calculate soil moisture values (Li et al. 1992, Li et al. 2000, Stange et al. 2000). The modelled amount of rainfall reaching the soil surface is influenced by interception (depending on LAI and rainfall history). Furthermore, winter storage of precipitation as snow is considered. Newer versions also account for surface runoff during heavy rainfall events. Actual soil moisture is calculated using a cascade model with determination of evapotranspiration losses. The agricultural version of DNDC uses a linear function to describe the effect of soil moisture on decomposition, but the *Forest-DNDC* version uses a Weibull function, which is allowed to vary from 0–1. These values are combined with a temperature factor to modify decomposition.

In *ROMUL* the monthly weight of moisture of organic horizons and mineral top soil are used to determine the influence or moisture availability on decomposition. If these data are missing, then a statistical soil weather generator SCLISS (Bykhovets and Komarov 2002) is used to calculate the moisture content based on the total monthly precipitation and hydrological soil characteristics. The calculation of potential evapotranspiration is based on the model that allows one to evaluate potential evapotranspiration using air temperature only (Blaney and Criddle 1950). Moisture of organic horizons is a non-linear rate modifier for decomposition rates of aboveground litter portions. Corresponding dependencies are evaluated from various experiments (Chertov et al. 2001). For belowground litter portions (root litter), the moisture of mineral topsoil is used to modify decomposition rates.

In *RothC* total monthly rainfall (mm) and monthly open pan evaporation (mm) (or PET, mm; (Smith et al. 2005b, Smith et al. 2006) are used to calculate topsoil moisture deficit (TSMD), as it is easier to obtain rainfall and pan evaporation data, from which the TSMD is calculated, than monthly measurements of the actual topsoil water deficit. Decomposition constants of active compartments are multiplied by a rate modifier that depends on cumulative monthly TSMD using a linear function within the typical range of TSMD. Outside the typical range, fixed minimum and maximum values are used.

In *Century*, soil moisture availability is simulated based on input data for rainfall, snow and irrigation, after adjusting for interception within the canopy, and taking into account water losses from simulated storm runoff, groundwater flow, and evapotranspiration (Metherell et al. 1993). Decomposition rates in *Century* are influenced by moisture availability according to a calculation based on the moisture left over from the previous monthly time step and current rainfall divided by PET. This estimation is combined with the temperature influence on decomposition to form a single multiplicative factor (Parton et al. 1994). The factor ranges from 0 to 1, with simulated values below 1 reducing decomposition rates.

In *Yasso*, the effects of moisture on decomposition are included in the same regression models that describe the effect of temperature on decomposition (see above) (Liski et al. 2003). These models are used to adjust the decomposition rates of model compartments. Moisture is estimated from a summer drought index calculated by subtracting potential evapotranspiration from precipitation.

2.2.3 Soil Texture

Soil texture has two main impacts on decomposition in the *ROMUL*, *RothC*, *Forest-DNDC* and *CENTURY* models. Texture influences decomposition both by affecting soil moisture through soil water holding capacity and by affecting stabilization of soil organic matter at higher clay contents.

SOILN model doesn't use texture as an input parameter; instead the hydraulic properties like

water retention curves, hydraulic conductivity and porosity are explicitly defined for each soil layer in the water and heat model *CoupModel*. The relationships between hydraulic properties and simulated soil moisture content, water flow and soil temperature are described in the *CoupModel* documentation (Jansson and Karlberg 2002). *SOILN* does not take into account the stabilizing effect of soil texture on SOM.

Unlike the other models, *Yasso* does not account for the effects of texture on decomposition. In *Yasso*, it is assumed that most of the short-term changes occur in organic layer where texture is not important, and that the effects of soil texture on soil C dynamics are mostly the result of influences on productivity litter production, rather than from the direct effect on decomposition (Liski et al. 2005).

2.2.4 Nitrogen

The group of the models selected in this study is very diverse with respect to nitrogen modeling. *SOILN*, *Forest-DNDC*, *ROMUL* and *CENTURY* explicitly include nitrogen in their framework coupled with soil C dynamics. These models require nitrogen input data such as atmospheric N deposition and N additions in fertilizers (Table 1). By contrast, nitrogen is ignored in *Yasso* and *RothC*.

In *SOILN*, plant N uptake is controlled by the plant C:N ratios and plant biomass, and growth is limited by N availability. N availability depends on litter quality, decomposition rates, immobilization and external inputs of nitrogen from fertilization, atmospheric deposition and nitrogen fixation. In addition, processes are included that determine losses of N, such as nitrate leaching, and gas emissions of CO₂ and NO₂ (Johnsson et al. 1987, Eckersten and Beier 1998, Eckersten et al. 1998).

In *Forest-DNDC* decomposition of SOM leads to the formation of inorganic ammonium, which can be processed by microbial activity into various reactive forms (e.g., NO₃⁻, NO₂⁻, N₂O, NO) and also to N₂, volatilized as NH₃ (depended on soil pH), chemo-denitrified (NO₂⁻)(depended on soil pH) or leached (depended on soil water flux, texture and plant N uptake). The algorithms

used for simulating soil N transformations are described in Li et al. (2000).

In *ROMUL*, the various litter cohorts and soil organic matter pools have different nitrogen contents, which govern the rates of their decomposition. The equations for nitrogen, being a principal limiting factor of plant nutrition in forest ecosystems, are similar to the equations for carbon, but have some additional kinetic parameters, reflecting nitrogen retention in the soil system (Chertov 1990). The total amount of nitrogen available for plants is determined at the ecosystem level and defines the forest growth. Nitrogen leaching and deposition, as well as gas emissions of CO₂ are included.

In *Century*, carbon uptake is simulated through plant production, and this process is limited by nitrogen availability according to a maximum C:N ratio (Metherell et al. 1993). Thus, nitrogen availability affects the amount of nitrogen taken up by the plant and its subsequent litter quality, as well as decomposition. In turn, nitrogen availability is affected by the rate of decomposition, the amount of nitrogen immobilized, and the external inputs of nitrogen to the model, including atmospheric deposition, biological nitrogen fixation and fertilization events. Parton et al. (1987) and Metherell et al. (1993) provide additional details.

Nitrogen is omitted in *RothC*, but is included in the Rothamsted Nitrogen Model (Bradbury et al. 1993), which has been further developed into the SUNDIAL model (Smith et al. 1996, Smith 2002) and currently being extended for use in organic soils as the ECOSSE model (Smith et al. 2005a).

2.2.5 Model Initialization

For each time step, the values for the state variables of any process model are dependent on parameterization of the processes, inputs to the model, and previous values of the state variables. Consequently, initialisation of model pools is a potentially critical step during the parameterization of a process-based model. The overall importance of initialisation will depend on the question being addressed, the time frame of the study, and the model structure.

The initialisation of model pools with measured

data is not straightforward, since model pools do not necessarily represent fractions measurable in the field (Table 4). Initialisation based upon measurements or on a subjective basis can lead to values for state variables that deviate considerably from those that would be achieved by running the model with any plausible combination of input parameters and data for estimating current trends. This can lead to simulations with excessively high loss or gain in carbon (in some models the trajectory can even be chaotic). In many model applications, this behaviour is avoided with a spin-up to a steady-state under native vegetation and mean climate, followed by simulations of the historical land use and management according to survey records and expert knowledge. The assumption of a soil being in a steady state equilibrium with respect to current inputs is likely to be violated in most applications; therefore, historical data is crucial to produce a more realistic initialisation of state variables, and quantify the current trends and short-term changes in soil carbon (Foster et al. 2003). Similarly, this assumption can be violated during the model calibration. The soil C pool of a calibration site(s) may be far from equilibrium, but still the model parameters are calibrated so that the modelled equilibrium state matches the measured C pool. Therefore, a further correction of the parameters and the pools is necessary to avoid the underestimation of century-term soil C accumulation (Wutzler and Reichstein 2007).

RothC can be initialized with spin-up runs and litter input data. In contrast to some of the other models, however, *RothC* can incorporate measurements in this initialisation process. For example, the size of the inert pool can be determined by ^{14}C measurements or using an almost linear relationship between the size of the inert pool and bulk soil carbon stocks (Falloon et al. 1998a). The remaining amount of the measured SOM forms the active pools in the model, which are assumed to be in steady state equilibrium. The *RothC* is run to steady state using an approximate litter input, and is then adjusted to achieve a perfect match with initial bulk soil C using the proportional difference between the measured carbon and that estimated using the first, approximate litter C inputs. The litter input corresponding to measurements (and past conditions) can be used to scale future litter input. *RothC* can also be run

in ‘reverse mode’ to deduct the required input corresponding to measured bulk soil content (Coleman et al. 1997). There are also promising results to determine initial sizes of all carbon pools of the Roth-C model from soil samples using soil fractionation (Zimmermann et al. 2006a) or infrared-spectroscopy (Zimmermann et al. 2006b). However, these procedures have only been tested for agricultural soils at this time.

The most common procedure for initialisation of the *Yasso* model is using spin-up runs with historical litter input. On national level analyses, *Yasso*’s initialization dominated the uncertainty of soil C balance but the use of longer spin-up period clearly reduced its effect (de Wit et al. 2006, Peltoniemi et al. 2006). Other options for the initialisation of the model are to allocate the measured values to model pools with some additional assumptions, or to calculate steady-state values analytically.

For *SOILN*, carbon or nitrogen for the different pools is often initialised by running the model several times for mature forests with stable fluxes (several spin-up runs). As a starting point for the spin-up runs, the sum of C and N of all pools in each layer equals the in-situ measured total sum of C and N. Total C and N is split into fine litter, coarse litter (i.e. woody litter), humus and microbes. Litter dominates in the surface layer(s), while in the mineral soil, litter is assumed to be proportional to root biomass distribution.

For the *Century* model, equations have been developed that provide a first approximation of initial pool sizes based on relationships derived from Burke et al. (1989). However, many applications begin with a spin-up to prevent chaotic behaviour.

The *Forest-DNDC* model is initialised with fixed proportions for each of the pools. These proportions will change over time based on litter input and its C:N ratio. Using spin-up runs for pool initialisation is also an alternative.

Although the *ROMUL* model can be initialised with spin-up runs and the steady state assumption, the model developers recommend using measured or compiled data on SOM and N pools in organic layer and mineral topsoil (Chertov, personal communication 2006). This is preferred because the pools correspond to soil layers that can be determined in the field. They have compiled a database

of initial values by forest site/type, tree species, age class, and area of consideration (Komarov and Chertov in press). Initial SOM and nitrogen pools of disturbed sites with unknown history can be initialised using forest successional series marked by vegetation composition.

2.3 Statistical Models of Soil Carbon

Another option for estimating changes in soil carbon stocks is to use common forest inventory variables and statistical methods. According to the UNFCCC, this is also considered a Tier 3 approach. An example of such a statistical approach to monitor soil carbon stocks is the U.S. *Forcarb* model, described by Smith and Heath (2002). Estimates for soil organic carbon are derived with statistical models from the STATSGO database and forest inventory data using the methodology described by Johnson and Kern (2002) and more recently by (Amichev and Galbraith 2004). Fundamentally, soil organic carbon stock changes at the national level are then estimated as a function of change in land cover and forest resources, including forest type and land use (US-EPA 2006).

At the regional scale, many parameters included in process-based soil models (e.g., soil moisture and temperature) co-vary with more general explanatory variables such as forest type, time since disturbance and location (Smith and Heath 2002). Therefore, this approach is likely to provide parallel trends as the process-based models.

3 Regional Soil Carbon Modeling

3.1 Overview on Data Sources

Information on land management, anthropogenic and natural disturbances are key inputs for the models. Small land-use changes may lead to large changes in soil carbon, e.g. land use change from cropland production to forest causes a relatively large increase in soil C stocks compared to many land use and management activities (Post and

Kwon 2000). Frequent disturbances induced by forest management (harvests, tillage), and natural disturbances (such as fire, pests) have an important effect on litter input subsequent years after the event (Gärdenäs 1998, Liski et al. 2002, Peltoniemi et al. 2004, Thürig et al. 2005). However, the direct effect of some disturbances on SOM is still controversial (Johnson and Curtis 2001).

National forest inventories (NFI) are carried in several countries over the world to provide representative data on e.g., forest (and other land) area, stand structure, soil type and timber resources (Tokola 2006). Together with land use surveys, and forest management records they provide the underlying input data on anthropogenic activity in forested lands (e.g. USDA 2005). Thus, they provide a valuable source of input data as well as a framework for scaling model results (Vetter et al. 2005, Perry et al. Submitted in 2006). Advantages of NFI data are that typically they cover a large area, in some cases the records span several decades. Temporal resolution, however, is often limited (1–10 years). Changes in the forests resulting from changes in growing conditions (e.g. climate change), harvesting or other disturbances, are measured but the information on the causes remains often limited. Inventory data (with appropriate biomass and biomass turnover models) constitute an representative nation-wide data source for independent soil modules, such as *Yasso* or *RothC*, that are not able to simulate plant production and litter input (see e.g. de Wit et al. 2006, see e.g. Liski et al. 2006, Smith et al. 2006).

Field measurements of forest inventories may not capture land-use changes or disturbances (thinnings, harvests, fire, etc) with sufficient precision and accuracy. Inventory grids are typically too coarse in comparison with what would be required to detect rare events occurring on limited area. Additional forest management records may be used to supplement NFI data, such as reported harvests or timber purchase statistics.

Remotely-sensed data provide spatial information to improve, for example, inventory estimates on growing stocks and land-use, or to produce information on vegetation cover to derive biomass estimates (Rosenqvist et al. 2003, Tomppo 2006). For example CORINE land cover data or the PELCOM database have been used as a basis for

model input data on land use and for up-scaling of measured soil carbon pools (e.g. Vleeshouwers and Verhagen 2002, Lettens et al. 2005a, Lettens et al. 2005b, Schmit et al. 2006) (Meesenburg et al. 1999). Digital elevation models (DEM) could also be used to provide information on topographic position, aspect and slope, which influence temperature and thermal regimes. In addition, edaphic characteristics are required by several models, including soil texture. The use of these data sources is important when plot level data are scaled over larger areas.

3.2 Scaling Model Functions and Input Data

Scaling decisions are a critical component of the inventory development. The resulting operational scale of the process-based model and associated input data will determine the spatio-temporal resolution of a model application, which could range from finely resolved scales such as m² and minutes to coarser scales such as 10s of km² and decades. In turn, the model application will determine the scale at which soil C stock trends can be interpreted. Interpreting trends at an inappropriate scale can introduce large biases, referred to as ecological fallacies (Clark 2003). Thus, scaling decisions will determine the level of detail available for reporting soil C trends as well as its uncertainty, which will be the basis for informing policy decisions.

Much of the experimental research used to generate models is conducted at relatively small spatial resolutions, such as a forest stand, and short time periods of weeks to a few years. Consequently, models developed from this information can be 1) applied repeatedly at a finer scale (e.g., a stand with daily time step), or 2) the model can be scaled to meet the spatio-temporal domain of the input data (Rastetter et al. 1992, Rastetter et al. 2003).

Repeated application of a model on finer scale often requires scaling of the input data to meet the operational scale of the model. For example, monthly mean temperature may need to be estimated for a daily time step. If scaling is not done, non-linear relationships between input and modeled output may cause biased estimates for the spatio-temporal domain of analysis.

Scaling models to a larger resolution can also create biases. For example, Ogle et al. (2006) demonstrated how increasing the spatial resolution of a model application by using coarser-scale parameters introduced significant bias into estimates of soil C change. Essentially, coarser scale parameters did not adequately represent the underlying regional heterogeneity in their analysis. Consequently, rescaling the model input or the model functions needs to be done with careful consideration of the uncertainties.

In many cases, input data sets are incomplete in time or space, for example, data may be collected only from a small number of plots for short period of time, which are subsequently scaled to a larger region. In these cases, interpolation and extrapolation can be used to estimate input values for missing data with techniques such as geostatistics (Kyriakidis 2001). Errors will inevitably be generated through these scaling exercises, and need to be quantified as part of the uncertainty in model results.

All models are developed for a certain spatio-temporal domain, and subsequently they neglect processes operating on a coarser domain. Increasing the spatio-temporal domain of a model application (e.g., from forest stand to a watershed or larger region) will impose additional processes on a model analysis. Ecosystems are structured in a hierarchical manner and different processes tend to operate and shape the structural patterns found within each level of the hierarchy (O'Neill et al. 1986, Holling 1992). Large scale processes may be considered invariant at a finer scale and treated as a constant if the system is relatively stable (Wessman 1992). For example, erosional processes or leaching are not likely to create significant heterogeneity for soil C stocks in a forest stand, over a short time period, but they will have a significant effect at a larger spatial scale where soil or dissolved organic C is redistributed by wind or water from the original positions to depositional sites (Harden et al. 1999, Liu et al. 2003, Kortelainen et al. 2006). Therefore, results from the models included in this review will likely have additional uncertainties unless they have been further developed or linked with another model representing the larger scale processes.

3.3 Model Selection: Influence of Data Availability

Model selection and development is the first step when developing forest inventories for soil C; this process has been documented thoroughly by Ogle and Paustian (2005). While any of the models included in this review could potentially be used for a national scale analysis, selection is constrained by the availability of model input and evaluation data.

Input data provide detail on spatial heterogeneity in initial conditions as well as associated change over time. These data characterize abiotic and biotic conditions as well as anthropogenic activity, and are fundamental to simulate variation in ecosystem process rates in a large-scale analysis. Although some of the variables affecting litter and SOM dynamics (such as soil texture, proportion of rocks) can be treated as static on each site, their inclusion in a regional analysis is often viewed as critical to extend the application range of the model (McGill 1996).

Fully process-based modeling approaches for ecosystems (see e.g. Chen et al. 2000), which simulate plant production, microbial decomposition and associated processes to provide estimates of changes in all C pools [typical application approach for *Century*, Forests-DNDC, *ROMUL* with forest process-based model of growth and element cycling *EFIMOD* (Komarov et al. 2003), and *SOILN*] have a tendency to require more input data than if soil pools are only modeled. For the latter, soil models are used in connection with measured inventory data (typical approach for *RothC* and *Yasso*), and appropriate biomass and biomass turnover models. Ultimately, availability of data dictates the selection between these two approaches.

With detailed forest inventory data, statistical models, such as *Forcarb* in US, may be used to estimate change in soil C stocks. However, this method is not transferable to new conditions without a large measurement effort, and is not typically used for forecasting soil C stock changes because it is predicated on measurements from the past that may not represent future conditions.

3.4 Model Input Data Requirements and Sources of Data

Climatic data from weather monitoring stations are available in most countries. Assuming that microclimatic conditions in (sample of) forests can be estimated with sufficient precision and that the soil or air temperatures can be adequately estimated by the models, there should not be a large difference among the models. Therefore, there is not likely to be a limitation in model selection based on weather data requirements. The one exception, however, may be the very detailed information required for several soil layers in *SOILN* and the fine time resolution of the models *SOILN*, *Forest-DNDC*.

Input data on soil texture and nitrogen are more likely to become limitations for model use than climatic data. These data are typically collected in soil surveys and in experimental studies (which may also provide data on soil C). Experimental soil studies at plot level may not provide sufficient coverage, or a link to other input data required by the models.

Soil texture (e.g., clay content) is needed in *Century*, Forests-DNDC, *ROMUL*, and *RothC*, and it is most often collected in surveys. However, in forest soils, this property varies widely, and plot averages may consist of very few samples. The input soil data requirements are greatest for *SOILN*, including hydraulic properties of each soil layer, but a database of hydraulic properties is provided for various soil textural compositions and land uses (Jansson and Karlberg 2002). If data on soil texture is missing, quality or coverage is limited, or its effect on soil C dynamics is assumed to be negligible, *Yasso* or a similar model can be used that does not require these input data.

The other potential limitation is the availability of nitrogen input data. Surveys usually provide information on average N concentration or C:N ratio in the upper soil layers, which may be used as an input parameter to initialize model nitrogen pools (in *Century*, Forests-DNDC, *ROMUL*, *SOILN*). If N data are not directly available, correlation between site fertility and N could allow an approximate estimation of initial N status of soil (Komarov and Chertov in press). The predictions of C and N with *Century*, Forests-DNDC,

ROMUL, *SOILN* also require information on nitrogen inputs such as atmospheric deposition and fertilization. Nitrogen fertilization is less common in forest, and data may be available from timber companies to the extent that fertilizers are used. Atmospheric N deposition is reported as NO_3^- and NH_4^+ , and these data are typically collected on large scales that could be used as inputs to the models requiring this information (see e.g., on-line data set from Holland et al. 2005).

4 Soil Model Evaluation

Besides availability of input data, the performance of the model can affect the model selection (Ogle and Paustian 2005). Evaluation data are used to test a model's performance, which along with an assessment of composition and sensitivity determine the adequacy of a model for reporting C stock changes (Prisley and Mortimer 2004). Specifically, the performance of soil models is evaluated by comparing simulation results to field measurements of stocks (see e.g. Smith et al. 1997) or fluxes of carbon.

An example of soil model evaluation for regional application is given by (Izaurrealde et al. 2001). They selected the model that performed best among six candidates by comparing model simulations to measurements in 7 long-term experiments. They used the selected model for a regional analysis of SOC changes in agricultural soils for two ecodistricts in Alberta, Canada. Besides, for model selection, evaluation of model performance can also be included as a component of the uncertainty, providing information on model error (Falloon and Smith 2003, Ogle et al. In press).

4.1 Comparability of Simulations and Empirical Data

Soil models vary in their definitions of carbon pools (Table 3) and the depth in the profile that is included in the model estimation of carbon stocks (Table 4). Similarly, soil inventories apply no consistent definitions for sampled soil pools, layers, and depth (Table 5). Testing model predictions

will clearly be influenced by varying definitions of measurable depth in soil inventories, which limits the usefulness of data for comparisons with model results (Table 5 and Table 6). Some of the models (*ROMUL*, *SOILN*, *RothC*, *Forest-DNDC*) facilitate comparisons to data of varying depths by allowing for an adjustable simulation depth during parameterization (Table 3).

Simulating distinctive pools of litter, an organic layer, and mineral soil layers could allow for more detailed comparisons with measurements. Some models (*Forest-DNDC*, *ROMUL*, *SOILN*) split decomposing matter to individual pools (litter, organic, and mineral), which also loosely correspond with vertical distribution of SOM. A newer model which was developed from *Century* also includes a separate humified litter pool (Nalder and Wein 2006). In *Forest-DNDC*, distinction was made due to its importance for soil hydrology, aeration and mineralization: the organic layer is simulated separately from the mineral soil (Li et al., 2000). In *ROMUL* and *Forest-DNDC*, the simulated organic layer pools are equivalent to a measurable pool. Simulating separate pools that are measurable for the litter and organic layer are especially relevant in boreal forests where the forest floor is distinctive and thick. Moreover, turnover rates of organic matter decrease with depth, which means that litter and organic layer contribute the most to the short term changes (Gaudinski et al. 2000). The changes of C in different soil layers may be even in opposite directions (e.g. Bashkin and Binkley 1998, Heidmann et al. 2002, Vesterdal et al. 2002). Even with a model that simulates measurable pools, comparing model results with existing measured data is often complicated because definitions of pools are not consistent, or are not even reported. For instance, among the soil inventories referred in this study, only Bellamy et al. (2005) explicitly mention that they separated the litter from the organic layer.

Attempts have also been made to characterize kinetic pools of SOM by chemical (Berg et al. 1991, Henriksen and Breland 1999), physical fractionation (Oades 1989, Elliot and Cambardella 1991, Cambardella and Elliot 1993, Oades 1993), and infrared spectroscopy (Zimmermann et al. 2006b, Zimmermann et al. 2006a). If these methods prove to be useful in future, they will most

Table 3. List of pools in soil models.

	<i>Yasso</i>	<i>ROMUL</i>	<i>SOILN</i>	<i>RothC</i>	<i>Forest-DNDC</i>	<i>CENTURY</i>
Stand	–	–	Plant: roots, stem, leaves (current year and older) and grains	–	3 vegetation layers (upper-story, understory, ground growth)	Plant production submodel with leaves, fine roots, fine branches, large wood, and coarse roots
Litter	2 (fine woody litter and coarse woody litter)	2 or more: Pools for both above-ground and belowground litter for different compartments of different tree species. Division of senescent plant material into litter pools based on nitrogen and ash contents	1 or 2 per soil layer, 10 to 15 soil layers	2 [resistant and decomposable plant material; RPM & DPM (Coleman and Jenkinson 1996)]	3 per soil layer (very labile litter, labile litter and resistant litter) per layer (Li et al. 2000)	Metabolic and structural pools for both aboveground and belowground litter. Division of senescent plant material into litter pools based on lignin/N ratio.
SOM	5 (extractives, celluloses, lignin-like compounds, 2 humus)	6 (or more if litter specified for different tree species)	1 humus pool and optional 1 microbe pool per soil layer (10–15 layers)	3 SOM pools: BIO, HUM & IOM	2 humads and humus per layer	3 SOM pools: active, slow and passive SOM
Pools divided between organic and mineral soil layers?	No	Yes	Yes	No	Yes, 2 layers in organic soil and 2 in mineral soil	No

likely be model-specific. This may limit their utility in evaluating several models as potential candidates for a regional analysis. Fractionation techniques may also be useful for model parameterization (Skjemstad et al. 2004), and initialisation of carbon pools (see also section 1.2.5).

Overall, soil surveys should provide high quality data and include error assessments of measured stocks and changes (Falloon and Smith 2003). Thus, ideally, all parameters needed to determine soil carbon stocks (C concentration, bulk density and stone content) should be measured to a sufficient soil depth, including the organic layer at forest sites, and with enough repetitions per unit area to represent the spatial variability of soil carbon stocks. As can be seen from in Table 5 and Table 6, parameters like bulk density or stone content are often not directly determined when

soil samples have been taken in the past. As a consequence, changes in soil carbon are often only expressed as changes in C concentrations (e.g. Bellamy et al. 2005, Kelly and Mays 2005).

Since models usually predict total amounts of carbon in the soil, C concentrations first have to be converted into stocks before they can be applied for model evaluation. This can be done by using various available pedotransfer functions to estimate bulk densities, which are often based on parameters such as soil texture (usually also only estimated in the field) and C concentrations (Tamminen and Starr 1994). This technique has been applied in a number of large-scale studies on soil C changes (e.g. Heidmann et al. 2002, Sleutel et al. 2003). The stone content can only be determined in the field although direct measurements are difficult and estimates often erroneous

Table 4. Specific model assumptions, limitations and planned improvements.

	<i>Yasso</i>	<i>ROMUL</i>	<i>SOILN</i>	<i>RothC</i>	<i>Forest-DNDC</i>	<i>CENTURY</i>
Simulation depth	Organic layer + 1 m mineral soil	Organic layer and 1 m mineral soils or the depth of measured data	Any depth, commonly 1.5–3 meters	Any depth to 1m	Routinely 50 cm, but can be modified to go down to e.g. 1.5 m	20 cm
Limitations	Only for upland forest soils	Only for well or excessively drained mineral soils	Needs a lot of input information, never been applied to peat soils	Only for upland forest soils	Needs a lot of input information	Only for top 20 cm, no peatlands, does not separate the humified portion of the litter from mineral soils (Kelly et al. 1997)
Planned improvements	–	Model for organic soils under development	<i>CoupModel</i> : new plant pool coarse roots, new harvest regimes modules	Model for organic soils under development (Smith et al. 2005a)	Version for Mediterranean forest under development	New model using <i>Century</i> -based algorithms with humified litter pool separate from mineral soil (Nalder and Wein 2006)
Measurability of pools	Only extractives, celluloses and sums of other pools measurable.	Yes, pools or their sums are measurable	No, conceptual pool approach which cannot directly be measured. Total sum of N and C can be compared with measured ones.	Yes (Skjemstad et al. 2004, Shirato and Yokozawa 2006, Zimmermann et al. 2006a)	No, conceptual pool approach, cannot directly be measured	Comparison of modeled SOC and measured C in SOM, as well as modeled and measured litter pools possible. Conceptual sub-pools for litter and soil organic C not directly measurable
Model application / evaluation	(Kaipainen et al. 2004, Peltoniemi et al. 2004, Palosuo et al. 2005, Thürig 2005, de Wit et al. 2006)	(Smith et al. 1997, Chertov et al. 1999, Chertov et al. 2002, Chertov et al. 2003, Mikhailov et al. 2004, Chertov et al. 2006)	Forest sites (Eckersten et al. 1995, Eckersten and Beier 1998, Beier et al. 2001, Gärdenäs et al. 2003). Evaluation (Tiktak and van Grinsven 1995, Wolf et al. 1996).	Forest sites (Coleman et al. 1997, Smith et al. 1997, Romanya et al. 2000, Thürig et al. 2006). Regional / continental application (Falloon et al. 1998b, Falloon et al. 2002, Smith et al. 2006).	Upland forest soils (Li et al. 2000, Stange et al. 2000, Butterbach-Bahl et al. 2001). Update for forested wetlands (Zhang et al. 2002, Cui et al. 2005). Tropical forests (Kiese et al. 2005). Model evaluation (Kesik et al. 2005, Miehle et al. 2006).	(Kelly et al. 1997, Smith et al. 1997, Peng et al. 1998)
Uncertainty or sensitivity analysis	(Liski et al. 2005, Peltoniemi et al. 2006, Monni et al. 2007)	(Grabarnik et al. 2007, Komarov and Chertov in press)	(Eckersten et al. 2001)	In unpublished reports – available from P. Smith	Most sensitive factor method or Monte Carlo approach (Li et al. 2004), applications (Kesik et al. 2005, Kiese et al. 2005)	None have been published.
Transferability of model to new country	Yes (Palosuo et al. 2005, Thürig et al. 2005, de Wit et al. 2006)	Yes (Nadporozhskaya et al. 2006, Shaw et al. 2006)	Yes, see (Eckersten et al. 1995, Eckersten and Beier 1998)	Yes, applied regionally and globally (Post et al. 1982, Jenkinson et al. 1991, Wang and Polglase 1995, Falloon et al. 1998a, Tate et al. 2000, Falloon and Smith 2002, Smith et al. 2005b, Smith et al. 2006)	Yes (Stange et al. 2000, Zhang et al. 2002, Kesik et al. 2005, Kiese et al. 2005)	Yes

Table 5. Repeated soil inventories at regional or country scale and information on data quality. (St: sampling time, Sd: soil depth sampled, Snp: Sample number per Plot, Pn: plot number, Pr: paired resampling, Bd: bulk density, Sc: stone content, Si: soil information as concentration only [gC kg⁻¹] or per area [C ha⁻¹]).

Country/ region	Land use	St	Sd [m]	Snp	Pn	Pr	Bd, Sc	Si
Great Britain ¹ England, Wales	forest, grassland cropland	1978, 1994, 1995, 2003*	0.15	25+ per 400 m ²	5662	Y	n	gC kg ⁻¹
Denmark ²	cropland	1987, 1998	0.5	1	336	Y	e	tC ha ⁻¹
Belgium ³	forest, grassland cropland	1990, 2000#	<0.3->1.0	differs LSU	289 LSU	N	m, e	tC ha ⁻¹
Belgium ⁴	forest, grassland cropland	1960, 1990, 2000	ca. 0.3	differs per LSU	289 LSU	N	m,e	tC ha ⁻¹
Belgium ⁵ Flanders	cropland	1989–91, 1992–95, 1996–99	0.24 (1.0)	total of 210 000 samples	7 APU	N	e	tC ha ⁻¹
Belgium ⁶ West Flanders	cropland	1952,1990, 2003	ca. 0.3	6+ per 50 m ²	116	Y	m,e	tC ha ⁻¹
Ireland ⁷ southeastern part	grassland	1964,1996	0.1	15, 25 per 400 m ²	191, 220	N	n	gC kg ⁻¹
Finland southern part ⁸	forest	1965–1993	0.1 / 0.3	8–25	54	N	Sc m	gC kg ⁻¹
Sweden ⁹	forest	1983–1987, 1993–2002 2003–2012	0–0.05, 0.45–0.55, 0.60–0.65	min. soil layers 1, humus 1–5	23 100	Y	e,e	tC ha ⁻¹

¹ Bellamy et al. 2005, ² Heidmann et al. 2002, ³ Lettens et al. 2005a, ⁴ Lettens et al. 2005b, ⁵ Sleutel et al. 2003, ⁶ Sleutel et al. 2006,

⁷ Zhang & McGrath 2004, ⁸ Tamminen & Derome 2005, ⁹ (Olsson 1999, Olsson et al. 2007)

* dependant on land-use, + bulked for C analyses, # for forests only 2000 data available

e: estimated from pedotransfer functions; m: measured; n: no / not determined; y: yes

LSU: landscape units derived from soil and land-cover data; APU: agropedological units

(Eriksson and Holmgren 1996, Wirth et al. 2004). A rapid approach that is easy to apply in inventories is to measure the steel rod penetration depth at several points in a plot and convert it to average stone content with empirical equations (Viro 1952, Tamminen 1991). Since the stone content influences total carbon stocks in the soil by reducing the available fine-soil volume (Leifeld et al. 2005), its exclusion from analyses will lead to erroneous results, especially in stone-rich forest sites (Jia and Akiyama 2005).

4.2 Large-scale Datasets of Soil Carbon Changes

Traditionally, long-term soil C studies focused on agricultural sites, where the soil organic matter content was important as a measure for soil fertility. Many of these studies pre-date the global change debate. The need for monitoring soil C contents across land-uses and large spatial scales only became important during the last few decades with rising interest in quantifying terrestrial carbon sources and sinks (Houghton et al. 1983).

The existing field experiments evaluating changes in European soil C at regional or coun-

Table 6. Repeated forest soil inventories at plot/ecosystem scale (St: sampling time, Sds: soil depth sampled, Snp: Sample number per Plot, Pn: plot number, Bd: bulk density, Sc: stone content, Ol: organic layer, Si: soil information as concentration only [gC kg⁻¹] or per area [C ha⁻¹], C: at least average temperature and rainfall data available, L: data on annual small litterfall available).

Country/ region	Forest type	St	Ol	Sds [m]	Snp	Pn	Bd, Sc	Si	C	L
New Zealand ¹	<i>Pinus radiata</i> , replanted in 1995/6 and 2000	before and after harvest	ni	≤ 2.0	5–30+ per 400m ²	30	m	tC ha ⁻¹	ni	ni
Germany ²	<i>Fagus sylvatica</i> and <i>Picea abies</i>	1966–1995	y	0.5	3–6+	2–6	ni	tC ha ⁻¹	y	y#
United States ³	mixed oak	1976, 2002	n	0.5	1	11	ni	gC kg ⁻¹	ni	ni
Camp Branch Watershed	deciduous and pine, regrowth after differ- ent harvest techniques	before and ca. 16 years after harvest	y, n	0.3–1.0	varying	4 forest sites, varying plot numbers	m,e	tC ha ⁻¹	y/ni	ni∅
United States ⁵	mixed oak, mixed oak- maple, mixed oak- basswood	11 times between 2001–2003	n	0.2	8+ per 450 m ²	3 fertilized/ unfertilized per forest	N	gC kg ⁻¹	ni	y

¹ Oliver et al. 2004, ² Meesenburg et al. 1999, ³ Kelly & Mays 2005, ⁴ Johnson et al. 2002, ⁵ Waldrop et al. 2004

* dependant on land-use, + bulked for C analyses, # available from other studies in the same forests, ∅ data on biomass available
e: estimated from pedotransfer functions; m: measured; n: no; y: yes; ni: no information

try scales is scarce overall and forested areas are less represented in these studies than agricultural land. Currently, four regional or country scale studies with repeated measurements for changes in forest soil properties have been reported (Table 5). Large scale surveys are valuable for model evaluation because they provide measurements made with coherent methods over large spatial coverage, which may constitute notable climatic gradients or other variables of importance, which are included in the models.

Although sample numbers per plot are generally limited in these experiments, average changes in subsets of data may be significant allowing for meaningful model comparison to the measure-

ments. Generally, paired re-sampling is necessary to detect change, and thus allows smaller sets of data to be compared to simulations. Unpaired sampling with no control plot can lead to a large amount of unexplained variation due to differences in historical management and other factors influencing soil C stocks (Garten and Wullschleger 1999).

A regional study in southern Finland evaluated changes in soil properties based on repeated unpaired sampling of 54 plots during 12–28 years (Tamminen and Derome 2005). The changes in organic layer content (%) were assessed down to 30 cm but changes in carbon (%) were analysed only for organic layer, which decreases

the usability of data in evaluation of models that cannot simulate organic layer explicitly. Changes in organic matter content of organic layer ($n_{\text{plots}} = 54$) and 0–30 cm mineral soil ($n_{\text{plots}} = 32$) were significant, as were the changes of N and C:N in organic layer.

A nationwide soil survey from Belgium (Letten et al. 2005a, Letten et al. 2005b) has been used to report soil C changes with unpaired repeated sampling. The change assessment was based on datasets of a number of individual soil surveys that represented specified landscape unit (LSU). For each LSU the average carbon content was then obtained from all soil profiles within this LSU and compared among different years of data collection (Letten et al. 2005a, Letten et al. 2005b). Significant changes in soil C in forested LSUs were rare due to large variability (Letten et al. 2005a).

In England and Wales, Bellamy et al. (2005) used a repeated paired soil sampling along a grid covering the entire region. However, only the top 15 cm was collected, and most of the land was not forested. Average of changes of soil C in groups of both coniferous and deciduous woodland were significant; group sizes were not reported (Bellamy et al. 2005).

Swedish soil survey provides a representative and repeated samples from a grid covering the whole nation (Olsson 1999, Ståhl et al. 2004). Measurements are conducted in conjunction of NFI, and they are concentrated on forest soils where stand growth exceeds $1 \text{ m}^3 \text{a}^{-1}$; soils under other land-uses are not measured. Third inventory cycle has started in 2003, and is scheduled to end in 2012 (Olsson, M. personal communication 2007).

4.3 Stand-scale Studies

Most studies evaluating changes in soil carbon are on the plot scale, and again forests are less represented than agricultural sites. The meta-database EuroSOMNET, (<http://www.rothamsted.bbsrc.ac.uk/aen/eusomnet/expts/eurodb.htm>) summarizes information about 110 long-term soil organic matter studies (Powlson et al. 1998, Smith et al. 2002) of which only two were located on forested land: a coniferous forest stand in Russia (started

in 1965) and a stand of natural forest regeneration in UK (started in 1883). These data have already been used for comparison with model results of *RothC* and *Century* (Falloon and Smith 2002).

Long-term studies in unmanaged old growth forests are scarce and based on a review of recently published studies, we are only aware of the long-term study at Hubbard Brook experimental forest (Fahey et al. 2005). Fahey et al. (2005) did not detect a significant change in SOM.

Other studies on forest soil carbon mainly focus on afforestation and on other disturbances like harvest, windthrow or fire followed by forest regrowth. It is likely that recently disturbed systems have the largest changes in soil C, and thus they may provide interesting test cases for models. Potential studies providing data for comparison have been summarized in recent literature (Johnson and Curtis 2001, Paul et al. 2002, Waldrop et al. 2004), while some additional experiments are included in Table 6.

4.4 Chronosequences in Model Evaluation

Since it can take decades before changes in soil carbon stocks are detectable (Johnson et al. 2002, Paul et al. 2002), chronosequence studies have often been used instead of time series data to assess management effects across time (Johnson and Curtis 2001). The assumption underlying chronosequence studies is that all sites are comparable and differences among sites are due to the parameter of interest (e.g., stand age, management). Since this assumption cannot be validated, chronosequence studies are associated with an unpredictable error due to site selection.

Chronosequences are likely to be valuable for model comparisons in future, providing that they are successful in isolating the variable of interest as the main determinant of differences between the sites. Johnson and Curtis (2001) and Paul et al. (2002) summarised a number of available chronosequence studies, and more recent studies analyzing the effects of afforestation and management on soil carbon stocks have been published since, including Vesterdal et al. (2002), Thuille and Schulze (2006), and Mund and Schulze (2006).

4.5 Future Prospects

A number of monitoring programs have started in recent years at various spatial scales and are now waiting for the first re-sampling. CarboEurope IP and NitroEurope IP may provide data for model comparison at the plot scale in the near future. A variety of model input parameters are being measured and intensive soil analyses conducted at six European forest sites (<http://www.carboeurope.org/>, <http://www.neu.ceh.ac.uk/>). Depending upon the extent, suitability and quality of plot level data (Table 3), these results may be used either for model calibration or testing. The combination of knowledge on the process and calibration in several representative sites from these studies may lead to enhanced predictive power of the models included in this review.

At the European scale, the soil inventory of the International Co-operative Programme (ICP) based on sampling with a grid density of 16 x 16 km² has been repeated after ten years following the first sampling. After processing samples collected in 2006, these results will provide a unique large-scale dataset that will likely be useful for model evaluation (BioSoil 2006).

5 Discussion

Development of forest carbon inventories is largely dependent on available data. We reviewed six process-based and one statistical model for estimating soil C stock changes, assessed how they incorporate input data, and listed references to studies that could potentially be used for model evaluation.

The models included in this review have been calibrated and applied in different regions, have been developed to simulate only certain ecosystem types, and require varying level of detail for input data (Table 1 and Table 4). Consequently, the methodology for applying these models in national GHG inventories will be somewhat unique. For example, models incorporating a module for ecosystem production may be used to prepare simultaneous estimates of biomass and litter production, which are dynamically linked to soil processes. Models restricted to simulation

of soil processes only, require input data on litter production.

The models with fewer requirements in respect to input data, such as *Yasso* or *RothC*, may be the only option for some countries given their resource availability. Models such as *Century*, *ROMUL*, *Forest-DNDC*, and *SOILN*, may be more accurate due to a greater level of detail represented in their structure. However, the input data requirements may be too great for their application. Regardless, it is recommended that inventory compilers consider and even test multiple models for which the necessary input data are available for their country.

Most of the models in this review are applicable on upland forest soils, while only the wetlands version of DNDC has been applied on peat lands (Zhang et al. 2002, Li et al. 2004). In the future, it is anticipated that versions of *RothC* and *ROMUL* will also be available for wetlands (Table 4). If model that is used in the soil C inventory is not applicable on peat soils, an alternative is to use IPCC Tier 1 method with default factors, or possibly the Tier 2 approach with habitat specific emission factors that are obtained with the flux measurements. However, use of same model across all land use categories, would increase the coherency of inventory. For example, change in land-use would not require changing the model, and re-calculating the time series (IPCC 2006).

At present, the number of published data sets that could be used for soil model evaluation is scarce (Table 5 and Table 6), but more measurements will become available in near future. Besides for measurements of soil carbon, evaluation of soil models requires comprehensive data on factors that influence SOM, as well as input and parameterization data. It seems that most of the existing studies provide sufficient information to evaluate only the simplest soil models without the need to use average or assumed values for some input variables. Hopefully, in future, more soil studies and inventories will collect and publish data on ancillary variables such as data on site history, stand description, climate, texture, and nitrogen inputs.

Process-based modelling of soil carbon dynamics can be used to produce soil C change estimates for countries with different resources and levels of input data, since a wide range of models with

differing input data requirements exists. Large uncertainties are involved in predicted soil C change estimates, comparisons of predictions between different countries, estimates prepared with different methodologies, and in comparisons to empirical data. Therefore, uncertainties of predicted soil C change estimates should be always assessed. Whenever possible, models and other methods should be developed in parallel.

Acknowledgements

This review is an offspring of an ESF-funded workshop devoted to assessment of current methodological development of large-scale forest carbon inventories was held at Koli, Finland in April 2006 (www.metla.fi/tapahtumat/2006/soil2006). We thank all the attendees of the workshop, especially the ones that were active in the discussions during the sessions, or had fun while gliding down the snowy hills of Koli Mountain.

Mikko Peltoniemi was funded by the European Commission through the Forest Focus project 'Monitoring changes in the carbon stock of forest soils' (www.metla.fi/hanke/843002) that was coordinated at Finnish Forest Research Institute.

References

Ågren, G.I. & Hyvönen, R. 2003. Changes in carbon stores in Swedish forest soils due to increased biomass harvest and increased temperatures analysed with a semi-empirical model. *Forest Ecology and Management* 174: 25–37.

Amichev, B.Y. & Galbraith, J.M. 2004. A revised methodology for estimation of forest soil carbon from spatial soils and forest inventory data sets. *Environmental Management* 33(1): 74–86.

Baldocchi, D.D. 2003. Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future. *Global Change Biology* 9(4): 479–492.

Bashkin, M.A. & Binkley, D. 1998. Changes in soil carbon following afforestation in Hawaii. *Ecology* 79(3): 828–833.

Beier, C., Eckersten, H. & Gundersen, P. 2001. Nitrogen cycling in a Norway spruce plantation in Denmark – A SOILN model application including organic N uptake. *The Scientific World* 1: 394–406.

Bellamy, P.H., Loveland, P.J., Bradley, R.I., Lark, R.M. & Kirk, G.J.D. 2005. Carbon losses from all soils across England and Wales 1978–2003. *Nature* 437(8): 245–248.

Berg, B. 2000. Litter decomposition and organic matter turnover in northern forest soils. *Forest Ecology and Management* 133(1–2): 13–22.

— & McClaugherty, C. 2003. *Plant litter: decomposition, humus formation, carbon sequestration*. Springer-Verlag. xii, 286 p. ISBN 3540443290.

—, Boeltink, H., Breymer, A., Ewertsson, A., Gallardo, A., Holm, B., Johansson, M.-B., Koivuoja, S., Meentemeyer, V., Nyman, P., Olofsson, J., Pettersson, A.-S., Reurslag, A., Staaf, H., Staaf, I. & Uba, L. 1991. Data on needle litter decomposition and soil climate as well as site characteristics for some coniferous forest sites. Part II. Decomposition data. Report 42, Swedish University of Agricultural Sciences, Department of Ecology and Environmental Research, Uppsala.

BioSoil 2006. ICP Forests' Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests. Part IIIa Sampling and analysis of soil. United nations economic commission for Europe convention on long-range transboundary air pollution and the Regulation of the European parliament and of the council concerning monitoring of forests and environmental interactions in the community (Forest Focus).

Blaney, H.F. & Criddle, W.D. 1950. Determining water requirements in irrigated areas from climatological and irrigation data. USDA Soil Cons. Serv. Techn. Paper 96.

Bradbury, N.J., Whitmore, A.P., Hart, P.B.S. & Jenkinson, D.S. 1993. Modeling the fate of nitrogen in crop and soil in the years following application of N-15-labeled fertilizer to winter-wheat. *Journal of Agricultural Science* 121: 363–379.

Burke, I.C., Yonker, C.M., Parton, W.J., Cole, C.V., Flach, K. & Schimel, D.S. 1989. Texture, climate, and cultivation effects on soil organic-matter content in US grassland soils. *Soil Science Society of America Journal* 53(3): 800–805.

Butterbach-Bahl, K., Stange, F., Papen, H. & Li, C.S. 2001. Regional inventory of nitric oxide and

- nitrous oxide emissions for forest soils of south-east Germany using the biogeochemical model PnET-N-DNDC. *Journal of Geophysical Research-Atmospheres* 106(D24): 34155–34166.
- Bykhovets, S.S. & Komarov, A.S. 2002. A simple statistical model of soil climate with a monthly step. *Eurasian Soil Science* 35(4): 392–400.
- Cambardella, C.A. & Elliot, E.T. 1993. Methods for physical separation and characterization of soil organic matter fractions. *Geoderma* 56: 449–457.
- Chen, J., Chen, W.J., Liu, J., Cihlar, J. & Gray, S. 2000. Annual carbon balance of Canada's forests during 1895–1996. *Global Biogeochemical Cycles* 14(3): 839–849.
- Chertov, O.G. 1990. SPECOM – A single tree model of pine stand / raw humus soil ecosystem. *Ecological Modelling* 50: 107–132.
- , Komarov, A.S. & Tsiplianovsky, A.M. 1999. The simulation of soil organic matter and nitrogen accumulation in Scots pine plantations on bare parent material using the combined forest model EFIMOD. *Plant and Soil* 213(1–2): 31–41.
- , Komarov, A.S., Nadporozhskaya, M., Bykhovets, S.S. & Zudin, S.L. 2001. ROMUL – a model of forest soil organic matter dynamics as a substantial tool for forest ecosystem modeling. *Ecological Modelling* 138: 289–308.
- , Komarov, A.S., Bykhovets, S.S. & Kobak, K.I. 2002. Simulated soil organic matter dynamics in forests of the Leningrad administrative area, north-western Russia. *Forest Ecology and Management* 169(1–2): 29–44.
- , Komarov, A., Kolström, M., Pitkänen, S., Strandman, H., Zudin, S. & Kellomäki, S. 2003. Modelling the long-term dynamics of populations and communities of trees in boreal forests based on competition for light and nitrogen. *Forest Ecology and Management* 176(1–3): 355–369.
- , Komarov, A., Loukianov, A., Mikhailov, A., Nadporozhskaya, M. & Zubkova, E. 2006. The use of forest ecosystem model EFIMOD for research and practical implementation at forest stand, local and regional levels. *Ecological Modelling* 194(1–3): 227–232.
- Christensen, B.T. 1996. Matching measurable soil organic matter fractions with conceptual pools in simulation models of carbon turnover: Revision of model structure. In: Powlson, D.S., Smith, P. & Smith, J.U. (eds.). *Evaluation of soil organic matter models*. NATO ASI Series I. 38. Springer-Verlag. p. 143–159.
- Clark, J.S. 2003. Uncertainty and variability in demography and population growth: a hierarchical approach. *Ecology* 84: 1370–1381.
- Coleman, K. & Jenkinson, D.S. 1996. RothC-26.3 – A Model for the turnover of carbon in soil. In: Powlson, D.S., Smith, P. & Smith, J.U. (eds.). *Evaluation of soil organic matter models – using existing long-term datasets*. NATO ASI Series Vol. I 38. Springer-Verlag, Heidelberg. p. 237–246.
- , Jenkinson, D.S., Crocker, G.J., Grace, P.R., Klir, J., Korschens, M., Poulton, P.R. & Richter, D.D. 1997. Simulating trends in soil organic carbon in long-term experiments using RothC-26.3. *Geoderma* 81(1–2): 29–44.
- Cui, J., Li, C., Sun, G. & Trettin, C.C. 2005. Linkage of MIKE SHE to Wetland-DNDC for carbon budgeting and anaerobic biogeochemistry simulation. *Biogeochemistry* (2005) 72: 147–167. *Biogeochemistry* 72: 147–167.
- Davidson, E.A. & Janssens, I.A. 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* 440: 165–173.
- , Trumbore, S.E. & Amundson, R. 2000. Biogeochemistry – soil warming and organic carbon content. *Nature* 408: 789–790.
- de Wit, H.A., Palosuo, T., Hylen, G. & Liski, J. 2006. A carbon budget of forest biomass and soils in southeast Norway calculated using a widely applicable method. *Forest Ecology and Management* 225(1–3): 15–26.
- Eckersten, H. & Beier, C. 1998. Comparison of N and C dynamics in two Norway spruce stands using a process oriented simulation model. *Environmental Pollution* 102: 395–401.
- , Gardenas, A. & Jansson, P.E. 1995. Modeling seasonal nitrogen, carbon, water and heat dynamics of the solling spruce stand. *Ecological Modelling* 83(1–2): 119–129.
- , Jansson, P.-E. & Johnsson, H. 1998. SOILN model, user's manual. Version 9.2. Swed. Univ. Agric. Sci., Dep. Soil Sci. Communications 98(6). 113 p.
- , Beier, C., Holmberg, M., Gundersen, P., Lepistö, A. & Persson, T. 1999. Application of the nitrogen model to forested land. In: *Nitrogen processes in arable and forest soils in the Nordic countries, field-scale modelling and experiments*. TemaNord 1999:560. Nordic Council of Ministers, Copenhagen. p. 99–129.

- 1 —, Blomback, K., Katterer, T. & Nyman, P. 2001.
2 Modelling C, N, water and heat dynamics in winter
3 wheat under climate change in southern Sweden.
4 *Agriculture Ecosystems & Environment* 86(3):
5 221–235.
- 6 Elliot, E.T. & Cambardella, C.A. 1991. Physical separa-
7 tion of soil organic matter. *Agriculture, Ecosystems
8 and Environment* 34: 407–419.
- 9 —, Paustian, K. & Frey, S.D. 1996. Modelling the
10 measurable or measuring the modelable: A hierar-
11 chical approach to isolating meaningful soil organic
12 matter fractionations. In: Powlson, D.S., Smith, P.
13 & Smith, J.U. (eds.). *Evaluation of soil organic
14 matter models*. NATO ASI Series I 38. Springer-
15 Verlag. p. 161–179.
- 16 Eriksson, C.P. & Holmgren, P. 1996. Estimating stone
17 and boulder content in forest soils – Evaluating the
18 potential of surface penetration methods. *Catena*
19 28(1–2): 121–134.
- 20 Fahey, T.J., Siccama, T.G., Driscoll, C.T., Likens, G.E.,
21 Campbell, J., Johnson, C.E., Battles, J.J., Aber,
22 J.D., Cole, J.J., Fisk, M.C., Groffman, P.M., Ham-
23 burg, S.P., Holmes, R.T., Schwarz, P.A. & Yanai,
24 R.D. 2005. The biogeochemistry of carbon at Hub-
25 bard Brook. *Biogeochemistry* 75(1): 109–176.
- 26 Falloon, P. & Smith, P. 2002. Simulating SOC changes
27 in long-term experiments with RothC and CEN-
28 TURY: model evaluation for a regional scale
29 application. *Soil Use and Management* 18(2):
30 101–111.
- 31 — & Smith, P. 2003. Accounting for changes in
32 soil carbon under the Kyoto Protocol: need for
33 improved long-term data sets to reduce uncertainty
34 in model projections. *Soil Use and Management*
35 19(3): 265–269.
- 36 —, Smith, P., Coleman, K. & Marshall, S. 1998a.
37 Estimating the size of the inert organic matter
38 pool from total soil organic carbon content for use
39 in the Rothamsted carbon model. *Soil Biology &
40 Biochemistry* 30(8–9): 1207–1211.
- 41 —, Smith, P., Smith, J.U., Szabo, J., Coleman, K. &
42 Marshall, S. 1998b. Regional estimates of carbon
43 sequestration potential: linking the Rothamsted
44 Carbon Model to GIS databases. *Biology and Fer-
45 tility of Soils* 27(3): 236–241.
- 46 —, Smith, P., Szabo, J. & Pasztor, L. 2002. Compari-
47 son of approaches for estimating carbon sequestra-
48 tion at the regional scale. *Soil Use and Management*
49 18: 164–174.
- 50 Foster, D., Swanson, F., Aber, J., Burke, I., Brokaw,
N., Tilman, D. & Knapp, A. 2003. The importance
of land-use legacies to ecology and conservation.
Bioscience 53: 77–88.
- Gärdenäs, A.I. 1998. Soil organic matter in European
forest floor in relation to stand characteristics and
environmental factors. *Scandinavian Journal of
Forest Research* 13: 274–283.
- , Eckersten, H. & Lillemägi, M. 2003. Modeling
long-term effects of N fertilization and N deposi-
tion on the N balance of forest stands in Sweden.
Emergo 2003(3). 30 p.
- Garten, C.T.J. & Wullschlegel, S.D. 1999. Soil carbon
inventories under a bioenergy crop (switchgrass):
measurement limitations. *Journal of Environmental
Quality* 28(4): 1359–1365.
- Gaudinski, J.B., Trumbore, S.E., Davidson, E.A. &
Zheng, S. 2000. Soil carbon cycling in a temperate
forest: radiocarbon-based estimates of residence
times, sequestration rates and partitioning fluxes.
Biogeochemistry 51: 33–69.
- Grabarnik, P., Bykhovets, S., Komarov, A., Mikhailov,
A. & Bezrukova, M. 2007. Global sensitivity analy-
sis of parameters of soil dynamic model ROMUL.
Manuscript.
- Harden, J.W., Sharpe, J.M., Parton, W.J., Ojima, D.S.,
Fries, T.I., Huntington, T.G. & Dabney, S.M. 1999.
Dynamic replacement and loss of soil carbon on
eroding cropland. *Global Biogeochemical Cycles*
13: 885–901.
- Heidmann, T., Christensen, B.T. & S.E., O. 2002.
Changes in soil C and N content in different crop-
ping systems and soil types. In: Petersen, S.O. &
Olesen, J.E. (eds.). *Greenhouse Gas Inventories
for Agriculture in the Nordic Countries Report*
81. Ministry of Food, Agriculture and Fisheries,
Danish Institute of Agricultural Sciences, Foulum.
p. 77–86.
- Henriksen, T.M. & Breland, T.A. 1999. Nitrogen avail-
ability effects on carbon mineralization, fungal
and bacterial growth, and enzyme activities during
decomposition of wheat straw in soil. *Soil Biology
& Biochemistry* 31: 1131–1134.
- Holland, E.A., Braswell, B.H., Sulzman, J.M. &
Lamarque, J.-F. 2005. Nitrogen deposition onto
the United States and Western Europe. Data set.
Available on-line [http://www.daac.ornl.gov] from
Oak Ridge National Laboratory Distributed Active
Archive Center, Oak Ridge, Tennessee, U.S.A.
- Holling, C.S. 1992. Cross-scale morphology, geometry,
and dynamics of ecosystems. *Ecological Mono-*

- graphs 64: 447–502.
- Houghton, R.A., Hobbie, J.E., Melillo, J.M., Moore, B., Peterson, B.J., Shaver, G.R. & Woodwell, G.M. 1983. Changes in the carbon content of terrestrial biota and soils between 1860 and 1980: a net release of CO₂ to the atmosphere. *Ecological Monographs* 53: 235–262.
- IPCC 2003. Good practice guidance for land use, land-use change and forestry. Institute for Global Environmental Strategies (IGES), Japan.
- Izaurrealde, R.C., Haugen-Kozyra, K.H., Jans, D.C., McGill, W.B., Grant, R.F. & Hiley, J.C. 2001. Soil C dynamics: Measurement, Simulation and site-to-region scale-up. In: Lal, R., Kimble, J.M., Follett, R.F. & Stewart, B.A. (eds.). *Assessment methods for soil carbon*. Lewis Publishers, New York, NY. p. 553–575.
- Jansson, P.-E. & Karlberg, L. 2002. COUP model – Coupled heat and mass transfer model for soil-plant-atmosphere system. Dept. of Civil and Environmental Engineering, Royal Institute of Technology. Available at <http://www.lwr.kth.se/Vara%20Datorprogram/CoupModel/index.htm>.
- Jenkinson, D.S., Adams, D.E. & Wild, A. 1991. Model estimates of CO₂ emissions from soil in response to global warming. *Nature* 351: 304–306.
- Jenny, H. 1941. *The factors of soil formation*. McGraw-Hill.
- Jia, S. & Akiyama, T. 2005. A precise, unified method for estimating carbon storage in cool-temperate deciduous forest ecosystems. *Agricultural and Forest Meteorology* 134(1–4): 70–80.
- Johnson, D.W. & Curtis, P.S. 2001. Effects of forest management on soil C and N storage: meta analysis. *Forest Ecology and Management* 140: 227–238.
- , Knoepp, J.D., Swank, W.T., Shan, J., Morris, L.A., Van Lear, D.H. & Kapeluck, P.R. 2002. Effects of forest management on soil carbon: results of some long-term resampling studies. *Environmental Pollution* 116: 201–208.
- Johnson, M.G. & Kern, J.S. 2002. Quantifying the organic carbon held in forested soils of the United States and Puerto Rico. In: Kimble, J.M., Heath, L.S., Birdsey, R.A. & Lal, R. (eds.). *The potential of U.S. forest soils to sequester and mitigate the greenhouse effect*, Lewis, Boca Raton. p. 47–72.
- Johnsson, H., Bergstrom, L., Jansson, P.E. & Paustian, K. 1987. Simulated nitrogen dynamics and losses in a layered agricultural soil. *Agriculture Ecosystems & Environment* 18(4): 333–356.
- Kaipainen, T., Liski, J., Pussinen, A. & Karjalainen, T. 2004. Managing carbon sinks by changing rotation length in European forests. *Environmental Science & Policy* 7(3): 205–219.
- Kätterer, T., Reichstein, M., Andrén, O. & Lomander, A. 1998. Temperature dependence of organic matter decomposition: a critical review using literature data analyzed with different models. *Biology and Fertility of Soils* 27: 258–262.
- , Eckersten, H., Heidmann, T., Borg, G., Sippola, J., Breland, T.A., Vold, A., Haugen, L.-E., Thomsen, A., Blombäck, K. & Bakken, L. 1999. Application of the nitrogen model to agricultural land. In: Jansson, P.-E., Persson, T. & Kätterer, T. (eds.). *Nitrogen processes in arable and forest soils in the Nordic Countries. – Field-scale modelling and experiments*. TemaNord 1999: 560. Nordic Council of Ministers, Copenhagen. p. 83–98.
- Kelly, J.M. & Mays, P.A. 2005. Soil carbon changes after 26 years in a Cumberland Plateau hardwood forest. *Soil Science Society of America Journal* 69(3): 691–694.
- Kelly, R.H., Parton, W.J., Crocker, G.J., Grace, P.R., Klir, J., Korschens, M., Poulton, P.R. & Richter, D.D. 1997. Simulating trends in soil organic carbon in long-term experiments using the century model. *Geoderma* 81(1–2): 75–90.
- Kesik, M., Ambus, P., Baritz, R., Bruggemann, N.B., Butterbach-Bahl, K., Damm, M., Duyzer, J., Horvath, L., Kiese, R., Kitzler, B., Leip, A., Li, C., Pihlatie, M., Pilegaard, K., Seufert, G., Simpson, D., Skiba, U., Smiatek, G., Vesala, T. & Zechmeister-Boltenstern, S. 2005. Inventories of N₂O and NO emissions from European forest soils. *Biogeosciences* 2(4): 353–375.
- Kiese, R., Li, C.S., Hilbert, D.W., Papen, H. & Butterbach-Bahl, K. 2005. Regional application of PnET-N-DNDC for estimating the N₂O source strength of tropical rainforests in the Wet Tropics of Australia. *Global Change Biology* 11(1): 128–144.
- Knorr, W., Prentice, I.C., House, J.I. & Holland, E.A. 2005. Long-term sensitivity of soil carbon turnover to warming. *Nature* 433: 298–301.
- Komarov, A. & Chertov, O. [Modelling of organic matter dynamics in forest ecosystems]. Nauka publishers, Moscow. 485 p. In press. (In Russian).
- , Chertov, O., Zudin, S., Nadporozhskaya, M., Mikhailov, A., Bykhovets, S., Zudina, E. & Zoubkova, E. 2003. EFIMOD 2 – a model of growth and cycling of elements in boreal forest ecosystems.

- Ecological Modelling 170: 373–392.
- Kortelainen, P., Rantakari, M., Huttunen, J.T., Mattsson, T., Alm, J., Juutinen, S., Larmola, T., Silvola, J. & Martikainen, P.J. 2006. Sediment respiration and lake trophic state are important predictors of large CO₂ evasion from small boreal lakes. *Global Change Biology* 12(8): 1554–1567.
- Kyriakidis, P.C. 2001. Geostatistical models of uncertainty for spatial data. In: Hunsaker, C.T., Goodchild, M.F., Friedl, M.A. & Case, T.J. (eds.). *Spatial uncertainty in ecology*. Springer-Verlag.
- Lagergren, F., Grelle, A., Lankreijer, H., Molder, M. & Lindroth, A. 2006. Current carbon balance of the forested area in Sweden and its sensitivity to global change as simulated by Biome-BGC. *Ecosystems* 9(6): 894–908.
- Leifeld, J., Bassin, S. & Fuhrer, J. 2005. Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. *Agriculture Ecosystems & Environment* 105(1–2): 255–266.
- Lettens, S., Van Orshoven, J., Van Wesemael, B., Muys, B. & Perrin, D. 2005a. Soil organic carbon changes in landscape units of Belgium between 1960 and 2000 with reference to 1990. *Global Change Biology* 11(12): 2128–2140.
- , Van Orshoven, J., van Wesemael, B., De Vos, B. & Muys, B. 2005b. Stocks and fluxes of soil organic carbon for landscape units in Belgium derived from heterogeneous data sets for 1990 and 2000. *Geoderma* 127(1–2): 11–23.
- Li, C., Cui, J., Sun, G. & Trettin, C.C. 2004. Modeling impacts of management on carbon sequestration and trace gas emissions in forested wetland ecosystems. *Environmental Management* 33(S1): S176–186.
- Li, C.S., Frolking, S. & Frolking, T.A. 1992. A model of nitrous-oxide evolution from soil driven by rainfall events. 2. Model applications. *Journal of Geophysical Research-Atmospheres* 97(D9): 9777–9783.
- , Aber, J., Stange, F., Butterbach-Bahl, K. & Papen, H. 2000. A process-oriented model of N₂O and NO emissions from forest soils: 1. Model development. *Journal of Geophysical Research-Atmospheres* 105(D4): 4369–4384.
- Liski, J., Perruchoud, D. & Karjalainen, T. 2002. Increasing carbon stocks in forest soils of western Europe. *Forest Ecology and Management* 169: 159–175.
- , Nissinen, A., Erhard, M. & Taskinen, O. 2003. Climatic effects on litter decomposition from arctic tundra to tropical rainforest. *Global Change Biology* 9(4): 575–584.
- , Palosuo, T., Peltoniemi, M. & Sievänen, R. 2005. Carbon and decomposition model Yasso for forest soils. *Ecological Modelling* 189: 168–182.
- , Lehtonen, A., Palosuo, T., Peltoniemi, M., Muukkonen, P., Eggers, T. & Mäkipää, R. 2006. Carbon accumulation in Finland's forests 1922–2004 – an estimate obtained by combination of forest inventory data with modelling of biomass, litter and soil. *Annales of Forest Science* 63(7): 687–697.
- Liu, S.R., Bliss, N., Sundquist, E. & Huntington, T.G. 2003. Modeling carbon dynamics in vegetation and soil under the impact of soil erosion and deposition. *Global Biogeochemical Cycles* 17: article 1074.
- Lloyd, J. & Taylor, J.A. 1994. On the temperature-dependence of soil respiration. *Functional Ecology* 8(3): 315–323.
- McGill, W.B. 1996. Review and classification of ten soil organic matter (SOM) models. In: Powlson, D.S., Smith, P. & Smith, J.U. (eds.). *Evaluation of soil organic matter models – using existing long-term datasets*. NATO ASI Series Vol. I 38. Springer-Verlag. p. 111–132.
- Meesenburg, H., Meiwes, K.-J. & Bartens, H. 1999. Veränderung der Elementvorräte im Boden von Buchen- und Fichtenökosystemen im Solling. *Berichte Freiburger Forstliche Forschung* 7: 109–114.
- Melillo, J.M., Aber, J.D. & Muratore, J.F. 1982. Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology* 63(3): 621–626.
- Metherell, A.K., Harding, L.A., Cole, C.V. & Parton, W.J. 1993. CENTURY: Soil organic matter model environment. GPSR Technical Report No. 4. United States Department of Agriculture, Fort Collins, Colorado, USA.
- Miehle, P., Livesley, S.J., Feikema, P.M., Li, C. & Arndt, S.K. 2006. Assessing productivity and carbon sequestration capacity of Eucalyptus globulus plantations using the process model forest-DNDC: Calibration and validation. *Ecological Modelling* 192(1–2): 83–94.
- Miglietta, F., Gioli, B., Hutjes, R.W.A. & Reichstein, M. 2007. Net regional ecosystem CO₂ exchange from airborne and ground-based eddy covariance, land-use maps and weather observations. *Global Change Biology* 13: 548–560.
- Mikhailov, A.V., Komarov, A.S. & Chertov, O.G. 2004.

- Simulation of the carbon budget for different scenarios of forest management. *Eurasian Soil Science* 37: 93–96.
- Monni, S., Peltoniemi, M., Palosuo, T., Mäkipää, R., Lehtonen, A. & Savolainen, I. 2007. Uncertainty of forest carbon stock changes – implications to the total uncertainty of GHG inventory of Finland. *Climatic Change* 81 (3–4): 391–413.
- Mund, M. & Schulze, E.D. 2006. Impacts of forest management on the carbon budget of European beech (*Fagus sylvatica*) forests. *Allgemeine Forst und Jagdzeitung* 177(3–4): 47–63.
- Nadporozhskaya, M.A., Mohren, G.M.J., Chertov, O.G., Komarov, A.S. & Mikhailov, A.V. 2006. Dynamics of soil organic matter in primary and secondary forest succession on sandy soils in The Netherlands: An application of the ROMUL model. *Ecological Modelling* 190(3–4): 399–418.
- Nalder, I.A. & Wein, R.W. 2006. A model for the investigation of long-term carbon dynamics in boreal forests of western Canada – I. Model development and validation. *Ecological Modelling* 192(1–2): 37–66.
- O'Neill, R.V., DeAngelis, D.L., Waide, J.B. & Allen, T.F.H. 1986. A hierarchical concept of ecosystems. Princeton University Press, Princeton NJ, USA.
- Oades, J.M. 1988. The retention of organic matter in soils. *Biogeochemistry* 5: 35–70.
- 1989. An introduction to organic matter in mineral soil. In: Dixon, J.B. & Weed, S.B. (eds.). *Minerals in soil environment*. Soil Science Society of America, Madison, WI, USA. p. 89–159.
- 1993. The role of biology in the formation, stabilization and degradation of soil structure. *Geoderma* 56: 377–400.
- Ogle, S.M., Breidt, F.J. & K., P. 2006. Bias and variance in model results associated with spatial scaling of measurements for parameterization in regional assessments. *Global Change Biology* 12: 516–523.
- & Paustian, K. 2005. Soil organic carbon as an indicator of environmental quality at the national scale: inventory monitoring methods and policy relevance. *Canadian Journal of Soil Science* 85: 531–540.
- , Breidt, F.J., Easter, M., Williams, S. & Paustian, K. Empirically based uncertainty associated with modeling carbon sequestration in soils. *Ecological Modelling* (in press).
- Olsson, M. 1999. Soil survey in Sweden. In: Bullock, P., Jones, R.J.A. & Montanarella, L. (eds.). *Soil resources of Europe*. Joint Research Centre, Ispra, Italy. p. 145–151.
- , Erlandsson, M., Lundin, L., Nilsson, T., Nilsson, Å. & Stendahl, J. 2007. Organic carbon in Swedish Podzols and their relation to site properties. *Biogeochemistry* (submitted).
- Palosuo, T., Liski, J., Trofymow, J.A. & Titus, B.D. 2005. Litter decomposition affected by climate and litter quality—Testing the Yasso model with litterbag data from the Canadian intersite decomposition experiment. *Ecological Modelling* 189(1–2): 183–198.
- Parton, W.J., Schimel, D.S., Cole, D.W. & Ojima, D.S. 1987. Analysis of factors controlling soil organic matter levels in Great Plains grasslands. *Soil Science Society of America Journal* 51: 1173–1179.
- , Ojima, D.S., Cole, C.V. & Schimel, D.S. 1994. A general model for soil organic matter dynamics: sensitivity to litter chemistry, texture and management. In: Bryant, R.B. & Arnold, R.W. (eds.). *Quantitative modeling of soil forming processes*. Soil Science Society of America, Special Publication 39, Madison, WI, USA. p. 147–167.
- Paul, K.I., Polglase, P.J., Nyakuengama, J.G. & Khanna, P.K. 2002. Change in soil carbon following afforestation. *Forest Ecology and Management* 168(1–3): 241–257.
- Peltoniemi, M., Mäkipää, R., Liski, J. & Tamminen, P. 2004. Changes in soil carbon with stand age – an evaluation of a modeling method with empirical data. *Global Change Biology* 10(12): 2078–2091.
- , Palosuo, T., Monni, S. & Mäkipää, R. 2006. Factors affecting the uncertainty of sinks and stocks of carbon in Finnish forests soils and vegetation. *Forest Ecology and Management* 232(1–3): 75–85.
- Peng, C., Apps, J.M., Price, D.T., Nalder, I.A. & Halliwell, D.H. 1998. Simulating carbon dynamics along the Boreal Forest Transect Case Study (BFTCS) in central Canada, 1, Model testing. *Global Biogeochemical Cycles* 12: 381–392.
- , Jiang, H., Apps, M.J. & Zhang, Y. 2002. Effects of harvesting regimes on carbon and nitrogen dynamics of boreal forests in central Canada: a process model simulation. *Ecological Modelling* 155: 177–189.
- Perry, C.H., Jandl, R., Peltoniemi, M., Matteucci, G. & Woodall, C. 2006. Detecting direct human-induced effects on the carbon balance of forest ecosystems

- 1 using models and national forest inventory data. 51
 2 (Submitted). 52
- 3 Post, W.M. & Kwon, K.C. 2000. Soil carbon sequestra- 53
 4 tion and land-use change: processes and potential. 54
 5 *Global Change Biology* 6: 317–327. 55
- 6 —, Emanuel, W.R., Zinke, P.J. & Stangenberger, 56
 7 A.G. 1982. Soil carbon pools and world life zones. 57
 8 *Nature* 298: 156–159. 58
- 9 —, Izaurralde, R.C., Mann, L.K. & Bliss, N. 2001. 59
 10 Monitoring and verifying changes of organic 60
 11 carbon in soil. *Climatic Change* 51(1): 73–99. 61
- 12 Powlson, D.S., Smith, P., Coleman, K., Smith, J.U., 62
 13 Glendining, M.J., Korschens, M. & Franko, U. 63
 14 1998. A European network of long-term sites 64
 15 for studies on soil organic matter. *Soil & Tillage 65*
 16 Research 47(3–4): 263–274. 66
- 17 Prescott, C.E. 1995. Does nitrogen availability control 67
 18 rates of litter decomposition in forests. *Plant and 68*
 19 *Soil* 169: 83–88. 69
- 20 Prisley, S.P. & Mortimer, M.J. 2004. A synthesis 70
 21 of literature on evaluation of models for policy 71
 22 applications, with implications for forest carbon 72
 23 accounting. *Forest Ecology and Management* 198: 73
 24 89–103. 74
- 25 Rastetter, E.B., King, A.W., Cosby, B.J., Hornberger, 75
 26 G.M., O'Neill, R.V. & Hobbie, J.E. 1992. Aggre- 76
 27 gating fine-scale ecological knowledge to model 77
 28 coarser-scale attributes of ecosystems. *Ecological 78*
 29 Applications 2: 55–70. 79
- 30 —, Aber, J.D., Peters, D.P.C., Ojima, D.S. & Burke, 80
 31 I.C. 2003. Using mechanistic models to scale eco- 81
 32 logical processes across space and time. *Bioscience* 82
 33 53: 68–76. 83
- 34 Ratkowsky, D.A., Olley, J., McMeekin, T.A. & Ball, 84
 35 A. 1982. Relationship between temperature and 85
 36 growth rate of bacterial cultures. *Journal of Bac- 86*
 37 teriology 149: 1–5. 87
- 38 Rodrigo, A., Recous, S., Neel, C. & Mary, B. 1997. 88
 39 Modelling temperature and moisture effects on 89
 40 C-N transformations in soils: comparison of nine 90
 41 models. *Ecological Modelling* 102(2–3): 325– 91
 42 339. 92
- 43 Rolff, C. & Ågren, G.I. 1999. A model study of nitro- 93
 44 gen limited forest growth. *Ecological Modelling* 94
 45 118: 193–211. 95
- 46 Romanya, J., Cortina, J., Falloon, P., Coleman, K. & 96
 47 Smith, P. 2000. Modelling changes in soil organic 97
 48 matter after planting fast-growing *Pinus radiata* on 98
 49 Mediterranean agricultural soils. *European Journal 99*
 50 of Soil Science 51(4): 627–641. 100
- Rosenqvist, A., Milne, A., Lucas, R., Imhoff, M. & 51
 Dobson, C. 2003. A review of remote sensing 52
 technology in support of the Kyoto Protocol. *Envi- 53*
 ronmental Science & Policy 6(5): 441–455. 54
- Schmit, C., Rousevell, M.D.A. & La Jeunesse, I. 2006. 55
 The limitations of spatial land use data in environ- 56
 mental analysis. *Environmental Science & Policy* 57
 9(2): 174–188. 58
- Shaw, C., Chertov, O.G., Komarov, A., Bhatti, J.S., 59
 Nadporozhskaya, M., Apps, J.M., Bykhovets, S. 60
 & Mikhailov, A. 2006. Application of the forest 61
 ecosystem model EFIMOD 2 to jack pine along 62
 the Boreal Forest Transect case study. *Canadian 63*
Journal of Soil Science 86: 171–185. 64
- Shirato, Y. & Yokozawa, M. 2006. Acid hydrolysis 65
 to partition plant material into decomposable 66
 and resistant fractions for use in the Rothamsted 67
 carbon model. *Soil Biology & Biochemistry* 38: 68
 812–816. 69
- Six, J., Callewaert, P., Lenders, S., de Gryze, S., Morris, 70
 S.J., Gregoriche, E.G., Paul, E.A. & Paustian, K. 71
 2002. Measuring and understanding carbon storage 72
 in afforested soils by physical fractionation. *Soil 73*
Science Society of America Journal. 74
- Skjemstad, J.O., Spouncer, L.R., Cowie, B. & Swift, 75
 R.S. 2004. Calibration of the Rothamsted organic 76
 carbon turnover model (RothC ver. 26.3), using 77
 measurable soil organic carbon pools. *Australian 78*
Journal of Soil Research 42(1): 79–88. 79
- Sleutel, S., De Neve, S., Hofman, G., Boeckx, P., 80
 Beheydt, D., Van Cleemput, O., Mestdagh, I., 81
 Lootens, P., Carlier, L., Van Camp, N., Verbeeck, 82
 H., Vande Walle, I., Samson, R., Lust, N. & Lemeur, 83
 R. 2003. Carbon stock changes and carbon seques- 84
 tration potential of Flemish cropland soils. *Global 85*
Change Biology 9(8): 1193–1203. 86
- Smith, J.E. & Heath, L.S. 2002. A model of forest floor 87
 carbon mass for United States forest types. Res. 88
 Pap. NE-722. USDA Forest Service, Northeastern 89
 Research Station, Newtown Square, PA. 37 p. 90
- Smith, J.U., Bradbury, N.J. & Addiscott, T.M. 1996. 91
 SUNDIAL: A PC-based system for simulating 92
 nitrogen dynamics in arable land. *Agronomy Jour- 93*
nal 88: 38–43. 94
- , Flynn, H.C. & Smith, P. 2005a. Development of 95
 the ECOSSE model. In: Flynn, H.C. & Smith, P. 96
 (eds.). Development of a model to simulate carbon 97
 and nitrogen dynamics in organic soils and predict 98
 response of these soils to land use and climate 99
 change, 2nd Annual Report on project UAB/013/03 100

- to SERRAD, Scottish Executive, Edinburgh, UK. p. 16–53.
- , Smith, P., Wattenbach, M., Zaehle, S., Hiederer, R., Jones, R.J.A., Montanarella, L., Rounsevell, M.D.A., Reginster, I. & Ewert, F. 2005b. Projected changes in mineral soil carbon of European croplands and grasslands, 1990–2080. *Global Change Biology* 11(12): 2141–2152.
- Smith, P. 2001. Soil organic matter modeling. In: Lal, R. (ed.) *Encyclopedia of soil science*. Marcel Dekker Inc, NY.
- 2002. Soil and the environment: Role of soil in models for climate change. In: Hillel, D., Rosenzweig, C., Powlson, D., Scow, K., Singer, M. & Sparks, D. (eds.). *Encyclopedia of soils in the environment*. Academic Press, London.
- , Smith, J.U., Powlson, D.S., McGill, W.B., Arah, J.R.M., Chertov, O.G., Coleman, K., Franko, U., Frokling, S., Jenkinson, D.S., Jensen, L.S., Kelly, R.H., Klein-Gunnewiek, H., Komarov, A.S., Li, C., Molina, J.A.E., Mueller, T., Parton, W.J., Thornley, J.H.M. & Whitmore, A.P. 1997. A comparison of the performance of nine soil organic matter models using datasets from seven long-term experiments. *Geoderma* 81: 153–225.
- , Falloon, P.D., Korschens, M., Shevtsova, L.K., Franko, U., Romanenkov, V., Coleman, K., Rodionova, V., Smith, J.U. & Schramm, G. 2002. EuroSOMNET – a European database of long-term experiments on soil organic matter: the WWW metadatabase. *Journal of Agricultural Science* 138: 123–134.
- , Smith, J.U., Wattenbach, M., Meyer, J., Lindner, M., Zaehle, S., Hiederer, R., Jones, R., Montanarella, L., Rounsevell, M., Reginster, I. & Kankaanpää, S. 2006. Projected changes in mineral soil carbon of European forests, 1990–2100. *Canadian Journal of Soil Science* 86: 159–169.
- Ståhl, G., Boström, B., Lindqvist, H., Lindroth, A., Nilsson, J. & Olsson, M. 2004. Methodological options for quantifying changes in carbon pools in Swedish forests. *Studia Forestalia Suecica* 214: 1–46.
- Stange, F. 2007. A novel approach to combine response functions in ecological process modelling. *Ecological Modelling* (in press).
- , Butterbach-Bahl, K., Papen, H., Zechmeister-Boltenstern, S., Li, C.S. & Aber, J. 2000. A process-oriented model of N₂O and NO emissions from forest soils 2. Sensitivity analysis and validation. *Journal of Geophysical Research-Atmospheres* 105(D4): 4385–4398.
- Sverdrup, H.U. 1990. The kinetics of base cation release due to chemical weathering. Lund University Press, Sweden. 246 p.
- , Warfvinge, P., Blake, L. & Goulding, K. 1995. Modeling recent and historic soil data from the Rothamsted-Experimental-Station, Uk Using Safe. *Agriculture Ecosystems & Environment* 53(2): 161–177.
- Tamminen, P. 1991. Kangasmaan ravinnetunnusten ilmaiseminen ja viljavuuden alueellinen vaihtelu Etelä-Suomessa. Summary: Expression of soil nutrient status and regional variation in soil fertility of forested sites in southern Finland. *Folia Forestalia* 777: 3–29.
- & Derome, J. 2005. Temporal trends in chemical parameters of upland forest soils in southern Finland. *Silva Fennica* 39(3): 313–330.
- & Starr, M. 1994. Bulk density of forested mineral soils. *Silva Fennica* 28(1): 53–60.
- Tate, K.R., Scott, N.A., Parshotam, A., Brown, L., Wilde, R.H., Giltrap, D.J., Trustring, N.A., Gomez, B. & Ross, D.J. 2000. A multi-scale analysis of a terrestrial carbon budget – Is New Zealand a source or sink of carbon? *Agriculture Ecosystems & Environment* 82(1–3): 229–246.
- Thuille, A. & Schulze, E.D. 2006. Carbon dynamics in successional and afforested spruce stands in Thuringia and the Alps. *Global Change Biology* 12(2): 325–342.
- Thürig, E. 2005. Carbon budget of Swiss forests: evaluation and application of empirical models for assessing future management impacts. PhD Thesis, Swiss Federal Institute of Technology, Zürich: p. 125.
- , Palosuo, T., Bucher, J. & Kaufmann, E. 2005. The impact of storm damage on carbon sequestration in Switzerland: a model-based assessment. *Forest Ecology and Management* 210(1–3): 337–350.
- , Faubert, P., Peltoniemi, M., Palosuo, T., Chertov, O.G., Komarov, A., Mikhailov, A., Suckow, F., Lasch, P., Wattenbach, M., Smith, P., Gottschalk, P. & Lindner, M. 2006. Comparison of soil carbon stocks and stock changes in forests simulated with four soil models. *Geoderma* (submitted manuscript).
- Tiktak, A. & van Grinsven, H.J.M. 1995. Review of sixteen forest-soil-atmosphere models. *Ecological Modelling* 83: 35–53.

- 1 Tokola, T. 2006. Part III Cases: Europe, Asia, North
2 America. In: Kangas, A. & Maltamo, M. (eds.).
3 Forest inventory, methodology and applications.
4 Managing Forest Ecosystems 10. Springer-Verlag.
5 p. 295–340.
- 6 Tomppo, E. 2006. The Finnish multi-source national
7 forest inventory – small area estimation and map
8 production. In: Kangas, A. & Maltamo, M. (eds.).
9 Forest inventory, methodology and applications.
10 Managing Forest Ecosystems 10. Springer-Verlag.
11 p. 195–224.
- 12 UNFCCC 1992. United Nations Framework Conven-
13 tion on Climate Change. [http://www.unfccc.de/
14 resource/convkp.html](http://www.unfccc.de/resource/convkp.html).
- 15 UNFCCC 1997. Kyoto Protocol. [http://www.unfccc.
16 de/resource](http://www.unfccc.de/resource).
- 17 UNFCCC 2006. National Inventory Submissions.
18 [http://unfccc.int/national_reports/annex_i_ghg_
19 inventories/national_inventories_submissions/
20 items/3734.php](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/3734.php).
- 21 US-EPA 2006. Inventory of U.S. greenhouse gas emis-
22 sions and sinks: 1990–2004. EPA 430-R-06-002.
23 U.S. Environmental Protection Agency, Washing-
24 ton, D.C. 459 p.
- 25 USDA 2005. Forest inventory and analysis national
26 core field guide. Vol. 1, Field Data Collection
27 Procedures for Phase 2 Plots, Ver. 3.0 [Online],
28 available at [www.fia.fs.fed.us/library/field-guides-
29 methods-proc](http://www.fia.fs.fed.us/library/field-guides-methods-proc) (verified 22 June 2006). USDA
30 Forest Service, Forest Inventory and Analysis
31 National Office, Arlington, Va.
- 32 Vesterdal, L., Ritter, E. & Gundersen, P. 2002. Change
33 in soil organic carbon following afforestation of
34 former arable land. *Forest Ecology and Manage-
35 ment* 169(1–2): 137–147.
- 36 Vetter, M., Wirth, C., Böttcher, H., Churkina, G.,
37 Schulze, E.-D., Wutzler, T. & Weber, G. 2005. Par-
38 titioning direct and indirect human-induced effects
39 on carbon sequestration of managed coniferous for-
40 ests using model simulations and forest inventories.
41 *Global Change Biology* 11(5): 810–827.
- 42 Viro, P.J. 1952. Kivisyyden määrittämisestä. Summary:
43 On the determination of stoniness. *Communicatio-
44 nes Instituti Forestalis Fenniae* 40(3). 23 p.
- 45 Vleeshouwers, L.M. & Verhagen, A. 2002. Carbon
46 emission and sequestration by agricultural land use:
47 a model study for Europe. *Global Change Biology*
48 8(6): 519–530.
- 49 Waldrop, M.P., Zak, D.R., Sinsabaugh, R.L., Gallo, M.
50 & Lauber, C. 2004. Nitrogen deposition modifies
soil carbon storage through changes in microbial
enzymatic activity. *Ecological Applications* 14(4):
1172–1177.
- Wang, Y.P. & Polglase, P.J. 1995. Carbon balance
in the tundra, boreal forest and humid tropical
forest during climate-change – scaling-up from
leaf physiology and soil carbon dynamics. *Plant
Cell and Environment* 18: 1226–1244.
- Wessman, C.A. 1992. Spatial scales and global change:
bridging the gap from plots to GCM grid cells.
Annual Review of Ecology and Systematics 23:
175–200.
- Wirth, C., Schulze, E.D., Schwalbe, G., Tomczyk, S.,
Weber, G. & Weller, E. 2004. Dynamik der Koh-
lenstoffvorräte in den Wäldern Thüringens. Series
Dynamik der Kohlenstoffvorräte in den Wäldern
Thüringens. Thüringer Landesanstalt für Wald,
Jagd und Fischerei, Gotha. 308 p.
- Wolf, J., Evans, L.G., Semenov, M.A., Eckersten, H. &
Iglesias, A. 1996. Comparison of wheat simulation
models under climate change. 1. Model calibration
and sensitivity analyses. *Climate Research* 7(3):
253–270.
- Wutzler, T. & Reichstein, M. 2007. Soils apart from
equilibrium – consequences for soil carbon balance
modelling. *Biogeosciences* 4: 125–136.
- Zhang, Y., Li, C.S., Trettin, C.C., Li, H. & Sun, G.
2002. An integrated model of soil, hydrology, and
vegetation for carbon dynamics in wetland ecosys-
tems. *Global Biogeochemical Cycles* 16(4).
- Zimmermann, M., Leifeld, J., Schmidt, M.W.I., Smith,
P. & Fuhrer, J. 2006a. Measured soil organic matter
fractions can be related to pools in the RothC
model. *European Journal of Soil Science* (submit-
ted).
- , Leifeld, J. & Fuhrer, J. 2006b. Quantifying soil
organic carbon fractions by infrared-spectroscopy.
Soil Biology and Biochemistry (submitted).

Total of 183 references